

CYCLIC M -CYCLE SYSTEMS OF COMPLETE GRAPHS MINUS A 1-FACTOR

HEATHER JORDON AND JOY MORRIS

ABSTRACT. In this paper, we provide necessary and sufficient conditions for the existence of a cyclic m -cycle system of $K_n - I$ when m and n are even and $m \mid n$.

In honor of Dan Archdeacon.

1. INTRODUCTION

Throughout this paper, K_n will denote the complete graph on n vertices, $K_n - I$ will denote the complete graph on n vertices with a 1-factor I removed (a *1-factor* is a 1-regular spanning subgraph), and C_m will denote the m -cycle (v_1, v_2, \dots, v_m) . An *m -cycle system* of a graph G is a set \mathcal{C} of m -cycles in G whose edges partition the edge set of G . An m -cycle system is called *hamiltonian* if $m = |V(G)|$.

Several obvious necessary conditions for an m -cycle system \mathcal{C} of a graph G to exist are immediate: $m \leq |V(G)|$, the degrees of the vertices of G must be even, and m must divide the number of edges in G . A survey on cycle systems is given in [4] and necessary and sufficient conditions for the existence of an m -cycle system of K_n and $K_n - I$ were given in [1, 17] where it was shown that an m -cycle system of K_n or $K_n - I$ exists if and only if $n \geq m$, every vertex of K_n or $K_n - I$ has even degree, and m divides the number of edges in K_n or $K_n - I$, respectively.

Throughout this paper, ρ will denote the permutation $(0 \ 1 \ \dots \ n - 1)$, so $\langle \rho \rangle = \mathbb{Z}_n$. An m -cycle system \mathcal{C} of a graph G with vertex set $V(G) = \mathbb{Z}_n$ is *cyclic* if, for every m -cycle $C = (v_1, v_2, \dots, v_m)$ in \mathcal{C} , the m -cycle $\rho(C) = (\rho(v_1), \rho(v_2), \dots, \rho(v_m))$ is also in \mathcal{C} . A cyclic n -cycle system \mathcal{C} of a graph G with vertex set \mathbb{Z}_n is called a *cyclic hamiltonian cycle system*. Finding necessary and sufficient conditions for cyclic m -cycle systems of K_n is an interesting problem and has attracted much attention (see, for example, [2, 3, 6, 7, 10, 11, 13, 15]). The obvious necessary conditions for a cyclic m -cycle system of K_n are the same as for an m -cycle system of K_n ; that is, $n \geq m \geq 3$, n is odd (so that the degree of every vertex is even), and m must divide the number of edges in K_n . However, these conditions are no longer necessarily sufficient. For example, it is not difficult to see that there is no cyclic decomposition of K_{15} into 15-cycles. Also, if p is an odd prime and $\alpha \geq 2$, then K_{p^α} cannot be decomposed cyclically into p^α -cycles [7].

The existence question for cyclic m -cycle systems of K_n has been completely settled in a few small cases, namely $m = 3$ [14], 5 and 7 [15]. For even m and $n \equiv 1 \pmod{2m}$, cyclic m -cycle systems of K_n are constructed for $m \equiv 0 \pmod{4}$ in [13] and for $m \equiv 2 \pmod{4}$ in [15]. Both of these cases are handled simultaneously in [10]. For odd m and $n \equiv 1$

(mod $2m$), cyclic m -cycle systems of K_n are found using different methods in [2, 6, 11]. In [3], as a consequence of a more general result, cyclic m -cycle systems of K_n for all positive integers m and $n \equiv 1 \pmod{2m}$ with $n \geq m \geq 3$ are given using similar methods. In [7], it is shown that a cyclic hamiltonian cycle system of K_n exists if and only if $n \neq 15$ and $n \notin \{p^\alpha \mid p \text{ is an odd prime and } \alpha \geq 2\}$. Thus, as a consequence of a result in [6], cyclic m -cycle systems of K_{2mk+m} exist for all $m \neq 15$ and $m \notin \{p^\alpha \mid p \text{ is an odd prime and } \alpha \geq 2\}$. In [18], the last remaining cases for cyclic m -cycle systems of K_{2mk+m} are settled, i.e., it is shown that, for $k \geq 1$, cyclic m -cycle systems of K_{2km+m} exist if $m = 15$ or $m \in \{p^\alpha \mid p \text{ is an odd prime and } \alpha \geq 2\}$. In [20], necessary and sufficient conditions for the existence of cyclic $2q$ -cycle and m -cycle systems of the complete graph are given when q is an odd prime power and $3 \leq m \leq 32$. In [5], cycle systems with a sharply vertex-transitive automorphism group that is not necessarily cyclic are investigated. As a result, it is shown in [5] that no cyclic m -cycle system of K_n exists if $m < n < 2m$ with n odd and $\gcd(m, n)$ a prime power. In [19], it is shown that if m is even and $n > 2m$, then there exists a cyclic m -cycle system of K_n if and only if the obvious necessary conditions that n is odd and that $n(n-1) \equiv 0 \pmod{2m}$ hold.

These questions can be extended to the case when n is even by considering the graph $K_n - I$. In [3], it is shown that for all integers $m \geq 3$ and $k \geq 1$, there exists a cyclic m -cycle system of $K_{2mk+2} - I$ if and only if $mk \equiv 0, 3 \pmod{4}$. In [12], it is shown that for an even integer $n \geq 4$, there exists a cyclic hamiltonian cycle system of $K_n - I$ if and only if $n \equiv 2, 4 \pmod{8}$ and $n \neq 2p^\alpha$ where p is an odd prime and $\alpha \geq 1$. In [8], it was shown that in every cyclic cycle decomposition of $K_{2n} - I$, the number of cycle orbits of odd length must have the same parity as $n(n-1)/2$. As a consequence of this result, in [8], it is shown that a cyclic m -cycle system of $K_{2n} - I$ can not exist if $n \equiv 2, 3 \pmod{4}$ and $m \not\equiv 0 \pmod{4}$ or $n \equiv 0, 1 \pmod{4}$ and m does not divide $n(n-1)$. In this paper we are interested in cyclic m -cycle systems of $K_n - I$ when m and n are even and $m \mid n$. The main result of this paper is the following.

Theorem 1.1. *For an even integer m and integer t , there exists a cyclic m -cycle system of $K_{mt} - I$ if and only if*

- (1) $t \equiv 0, 2 \pmod{4}$ when $m \equiv 0 \pmod{8}$,
- (2) $t \equiv 0, 1 \pmod{4}$ when $m \equiv 2 \pmod{8}$ with $t > 1$ if $m = 2p^\alpha$ for some prime p and integer $\alpha \geq 1$,
- (3) $t \geq 1$ when $m \equiv 4 \pmod{8}$, and
- (4) $t \equiv 0, 3 \pmod{4}$ when $m \equiv 6 \pmod{8}$.

Our methods involve circulant graphs and difference constructions. In Section 2, we give some basic definitions and lemmas while the proof of Theorem 1.1 is given in Sections 3, 4 and 5. In Section 3, we handle the case when $m \equiv 0 \pmod{8}$ and show that there is a cyclic m -cycle system of $K_{mt} - I$ if and only if $t \geq 2$ is even. In Section 4, we handle the case when $m \equiv 4 \pmod{8}$ and show that there is a cyclic m -cycle system of $K_{mt} - I$ if and only if $t \geq 1$. In Section 5, we handle the case when $m \equiv 2 \pmod{4}$. When $m \equiv 2 \pmod{8}$, we show that there is a cyclic m -cycle system of $K_{mt} - I$ if and only if $t \equiv 0, 1 \pmod{4}$. When $m \equiv 6 \pmod{8}$, we show that there is a cyclic m -cycle system of $K_{mt} - I$ if and only if $t \equiv 0, 3 \pmod{4}$. Our main theorem then follows.

2. PRELIMINARIES

The notation $[1, n]$ denotes the set $\{1, 2, \dots, n\}$. The proof of Theorem 1.1 uses circulant graphs, which we now define. For $x \not\equiv 0 \pmod{n}$, the *modulo n length* of an integer x , denoted $|x|_n$, is defined to be the smallest positive integer y such that $x \equiv y \pmod{n}$ or $x \equiv -y \pmod{n}$. Note that for any integer $x \not\equiv 0 \pmod{n}$, it follows that $|x|_n \in [1, \lfloor \frac{n}{2} \rfloor]$. If L is a set of modulo n lengths, we define the *circulant graph* $\langle L \rangle_n$ to be the graph with vertex set \mathbb{Z}_n and edge set $\{\{i, j\} \mid |i - j|_n \in L\}$. Notice that in order for a graph G to admit a cyclic m -cycle decomposition, G must be a circulant graph, so circulant graphs provide a natural setting in which to construct cyclic m -cycle decompositions.

The graph K_n is a circulant graph, since $K_n = \langle \{1, 2, \dots, \lfloor n/2 \rfloor\} \rangle_n$. For n even, $K_n - I$ is also a circulant graph, since $K_n - I = \langle \{1, 2, \dots, (n-2)/2\} \rangle_n$ (so the edges of the 1-factor I are of the form $\{i, i + n/2\}$ for $i = 0, 1, \dots, (n-2)/2$).

Let H be a subgraph of a circulant graph $\langle L \rangle_n$. The notation $\ell(H)$ will denote the set of modulo n edge lengths belonging to H , that is,

$$\ell(H) = \{\ell \in L \mid \{g, g + \ell\} \in E(H) \text{ for some } g \in \mathbb{Z}_n\}.$$

Many properties of $\ell(H)$ are independent of the choice of L ; in particular, the next lemma in this section does not depend on the choice of L .

Let C be an m -cycle in circulant graph $\langle L \rangle_n$ and recall that the permutation $\rho = (0\ 1\ \dots\ n-1)$, which generates \mathbb{Z}_n , has the property that $\rho(C) \in \mathcal{C}$ whenever $C \in \mathcal{C}$. We can therefore consider the action of \mathbb{Z}_n as a permutation group acting on the elements of \mathcal{C} . Viewing matters this way, the *length of the orbit of C* (under the action of \mathbb{Z}_n) can be defined as the least positive integer k such that $\rho^k(C) = C$. Observe that such a k exists since ρ has finite order; furthermore, the well-known orbit-stabilizer theorem (see, for example [9, Theorem 1.4A(iii)]) tells us that k divides n . Thus, if G is a graph with a cyclic m -cycle system \mathcal{C} with $C \in \mathcal{C}$ in an orbit of length k , then it must be that k divides $n = |V(G)|$ and that $\rho(C), \rho^2(C), \dots, \rho^{k-1}(C)$ are distinct m -cycles in \mathcal{C} .

The next lemma gives many useful properties of an m -cycle C in a cyclic m -cycle system \mathcal{C} of a graph G with $V(G) = \mathbb{Z}_n$ where C is in an orbit of length k . Many of these properties are also given in [7] in the case that $m = n$. The proofs of the following statements follow directly from the previous definitions and are therefore omitted.

Lemma 2.1. *Let \mathcal{C} be a cyclic m -cycle system of a graph G of order n and let $C \in \mathcal{C}$ be in an orbit of length k . Then*

- (1) $|\ell(C)| = mk/n$;
- (2) C has n/k edges of length ℓ for each $\ell \in \ell(C)$;
- (3) $(n/k) \mid \gcd(m, n)$;

Let $k > 1$ and let $P : v_0 = 0, v_1, \dots, v_{mk/n}$ be a subpath of C of length mk/n . Then

- (4) if there exists $\ell \in \ell(C)$ with $k \mid \ell$, then $m = n/\gcd(\ell, n)$,
- (5) $v_{mk/n} = kx$ for some integer x with $\gcd(x, n/k) = 1$,
- (6) $v_1, v_2, \dots, v_{mk/n}$ are distinct modulo k ,
- (7) $\ell(P) = \ell(C)$, and
- (8) $P, \rho^k(P), \rho^{2k}(P), \dots, \rho^{n-k}(P)$ are pairwise edge-disjoint subpaths of C .

Let X be a set of m -cycles in a graph G with vertex set \mathbb{Z}_n such that $\mathcal{C} = \{\rho^i(C) \mid C \in X, i = 0, 1, \dots, n-1\}$ is an m -cycle system of G . Then X is called a *generating set* for \mathcal{C} . Clearly, every cyclic m -cycle system \mathcal{C} of a graph G has a generating set X as we may always let $X = \mathcal{C}$. A generating set X is called a *minimum generating set* if $C \in X$ implies $\rho^i(C) \notin X$ for $1 \leq i \leq n$ unless $\rho^i(C) = C$.

Let \mathcal{C} be a cyclic m -cycle system of a graph G with $V(G) = \mathbb{Z}_n$. To find a minimum generating set X for \mathcal{C} , we start by adding C_1 to X if the length of the orbit of C_1 is maximum among the cycles in \mathcal{C} . Next, we add C_2 to X if the length of the orbit of C_2 is maximum among the cycles in $\mathcal{C} \setminus \{\rho^i(C_1) \mid 0 \leq i \leq n-1\}$. Continuing in this manner, we add C_3 to X if the length of the orbit of C_3 is maximum among the cycles in $\mathcal{C} \setminus \{\rho^i(C_1), \rho^i(C_2) \mid 0 \leq i \leq n-1\}$. We continue in this manner until $\{\rho^i(C) \mid C \in X, 0 \leq i \leq n-1\} = \mathcal{C}$. Therefore, every cyclic m -cycle system has a minimum starter set. Observe that if X is a minimum generating set for a cyclic m -cycle system \mathcal{C} of the graph $\langle L \rangle_n$, then it must be that the collection of sets $\{\ell(C) \mid C \in X\}$ forms a partition of L .

In this paper, we are interested in the cyclic m -cycle systems of $K_n - I$ where $n = mt$ for some positive integer t . Suppose K_n has a cyclic m -cycle system \mathcal{C} for some $n = mt$. Let X be a minimum generating set for \mathcal{C} and let $C \in X$ be a cycle in an orbit of length k . Then, $\ell(C)$ has $mk/n = k/t$ lengths which implies that $k = \ell t$ for some integer ℓ . Also, since $|\ell(C)| = \ell$, it follows that $\ell \mid m$. The following lemma will be useful in determining the congruence classes of t based on the congruence class of m modulo 8.

Lemma 2.2. *Let m be an even integer and let $K_{mt} - I$ have a cyclic m -cycle system for some positive integer t .*

- (1) *If $\{1, 2, \dots, (mt-2)/2\}$ has an odd number of even integers, then t is even.*
- (2) *If $\{1, 2, \dots, (mt-2)/2\}$ has an odd number of odd integers, then t is odd.*

Proof. Let m be even and suppose $K_{mt} - I$ has a cyclic m -cycle system \mathcal{C} for some positive integer t . Let $V(K_{mt}) = \mathbb{Z}_{mt}$, and let X be a minimum generating set for \mathcal{C} .

Suppose first that $\{1, 2, \dots, (mt-2)/2\}$ has an odd number of even integers. Since the set $\{\ell(C) \mid C \in X\}$ is a partition of $\{1, 2, \dots, (mt-2)/2\}$, there must be an odd number of cycles C in X with $\ell(C)$ containing an odd number of evens. Let $C \in X$ be a cycle in an orbit of length k with an odd number of even edge lengths. Let $|\ell(C)| = \ell$ and note that $k = \ell t$. From Lemma 2.1, we know that the subpath of C starting at vertex 0 of length ℓ ends at vertex jk with $\gcd(j, m/\ell) = 1$.

Suppose first k is odd. Then ℓ and t must both be odd. Thus m/ℓ is even so that jk is odd. Hence, $\ell(C)$ contains an odd number of odd integers and, since $|\ell(C)|$ is odd, an even number of even integers, contradicting the choice of C . Thus, k is even. Since k is even, jk is even. Thus, $\ell(C)$ contains an even number of odd integers. If ℓ is even, then $\ell(C)$ also contains an even number of even integers, contradicting the choice of C . Thus, ℓ is odd. Since k is even and $k = \ell t$, it must be that t is even.

Now suppose $\{1, 2, \dots, (mt-2)/2\}$ has an odd number of odd integers. Hence there are an odd number of cycles C in X with $\ell(C)$ containing an odd number of odd integers. Again, let $C \in X$ be such a cycle with $|\ell(C)| = \ell$, in an orbit of length $k = \ell t$. Let the subpath of C starting at vertex 0 of length ℓ end at vertex jk with $\gcd(j, m/\ell) = 1$. Now, if k is

even, then jk is even so that $\ell(C)$ contains an even number of odd integers, contradicting the choice of C . Thus k is odd. Since $k = \ell t$, we have that t is odd. \square

The following corollary is an immediate consequence of Lemma 2.2 and [12].

Corollary 2.3. *For an even integer m and a positive integer t , if there exists a cyclic m -cycle system of $K_{mt} - I$, then*

- (1) $t \equiv 0, 2 \pmod{4}$ when $m \equiv 0 \pmod{8}$,
- (2) $t \equiv 0, 1 \pmod{4}$ when $m \equiv 2 \pmod{8}$ with $t > 1$ if $m = 2p^\alpha$ for some prime p and integer $\alpha \geq 1$,
- (3) $t \equiv 0, 3 \pmod{4}$ when $m \equiv 6 \pmod{8}$, and
- (4) $t \geq 1$ when $m \equiv 4 \pmod{8}$.

Let $n > 0$ be an integer and suppose there exists an ordered m -tuple (d_1, d_2, \dots, d_m) satisfying each of the following:

- (i) d_i is an integer for $i = 1, 2, \dots, m$;
- (ii) $|d_i| \neq |d_j|$ for $1 \leq i < j \leq m$;
- (iii) $d_1 + d_2 + \dots + d_m \equiv 0 \pmod{n}$; and
- (iv) $d_1 + d_2 + \dots + d_r \not\equiv d_1 + d_2 + \dots + d_s \pmod{n}$ for $1 \leq r < s \leq m$.

Then an m -cycle C can be constructed from this m -tuple, that is, let $C = (0, d_1, d_1 + d_2, \dots, d_1 + d_2 + \dots + d_{m-1})$, and $\{C\}$ is a minimum generating set for a cyclic m -cycle system of $\langle \{d_1, d_2, \dots, d_m\} \rangle_n$. Thus, in what follows, to find cyclic m -cycle systems of $\langle L \rangle_n$, it suffices to partition L into m -tuples satisfying the above conditions. Hence, an m -tuple satisfying (i)-(iv) above is called a *difference m -tuple* and it *corresponds* to the m -cycle $C = (0, d_1, d_1 + d_2, \dots, d_1 + d_2 + \dots + d_{m-1})$ in $\langle L \rangle_n$.

3. THE CASE WHEN $m \equiv 0 \pmod{8}$

In this section, we consider the case when $m \equiv 0 \pmod{8}$ and show that there exists a cyclic m -cycle system of $K_{mt} - I$ for each even positive integer t . We begin with the case $t = 2$.

Lemma 3.1. *For each positive integer $m \equiv 0 \pmod{8}$, there exists a cyclic m -cycle system of $K_{2m} - I$.*

Proof. Let m be a positive integer such that $m \equiv 0 \pmod{8}$, say $m = 8r$ for some positive integer r . Then $K_{2m} - I = \langle S' \rangle_{2m}$ where $S' = \{1, 2, \dots, m-1\} = \{1, 2, \dots, 8r-1\}$. The proof proceeds as follows. We begin by finding a path P of length $m/2 = 4r$, ending at vertex m , so that $C = P \cup \rho^m(P)$ is an m -cycle. Note that $\langle \{2\} \rangle_{2m}$ consists of two vertex disjoint m -cycles. For the remaining $4r - 2$ edge lengths in $S' \setminus (\ell(P) \cup \{2\})$, we find $2r - 1$ paths P_i of length 2, ending at vertex 4 or -4 , so that $C_i = P_i \cup \rho^4(P_i) \cup \rho^8(P_i) \cup \dots \cup \rho^{2m-4}(P_i)$ is an m -cycle. Then this collection of cycles will give a minimum generating set for a cyclic m -cycle system of $K_{2m} - I$.

Suppose first that r is odd. For $r = 1$, let $P : 0, -3, 3, 7, 8$ and note that the edge lengths of P in the order encountered are 3, 6, 4, 1. For $r = 3$, let

$$P : 0, -3, 3, -7, 7, -11, 11, 23, 19, 20, -20, -4, 24$$

and note that edge lengths of P in the order encountered are 3, 6, 10, 14, 18, 22, 12, 4, 1, 8, 16, 20. For $r \geq 5$, let

$$P : 0, -3, 3, -7, 7, \dots, -(4r-1), 4r-1, 8r-1, 8r-5, 8r-4, 8r+4, 8r-8, 8r+8, \dots, \\ 6r+2, 10r-2, 6r-10, 10r+2, 6r-14, \dots, 12r-8, 4r-4, 8r$$

be a path of length $m/2$ whose edge lengths in the order encountered are 3, 6, 10, 14, \dots , $8r-6$, $8r-2$, $4r$, 4, 1, 8, 12, 16, \dots , $4r-4$, $4r+8$, $4r+12$, \dots , $8r-8$, $8r-4$, $4r+4$.

Now suppose that r is even. For $r=2$, let $P : 0, -3, 3, -7, 7, -1, -5, -4, 16$ and note that the edge lengths of P in the order encountered are 3, 6, 10, 14, 8, 4, 1, 12. For $r \geq 4$, let

$$P : 0, -3, 3, -7, 7, \dots, -(4r-1), 4r-1, -1, -5, -4, 4, -8, 8, \dots, \\ -(2r-4), 2r-4, -2r, 2r+8, -(2r+4), 2r+12, \dots, -(4r-8), 4r, -(4r-4), 8r$$

be a path of length $m/2$ whose edge lengths in the order encountered are 3, 6, 10, 14, \dots , $8r-6$, $8r-2$, $4r$, 4, 1, 8, 12, 16, \dots , $4r-8$, $4r-4$, $4r+8$, $4r+12$, \dots , $8r-8$, $8r-4$, $4r+4$.

In each case, let $C = P \cup \rho^m(P)$ and observe that C is an m -cycle C with $\ell(C) = \{1, 3, 4, 6, 8, \dots, 8r-2\}$. Let $C' = (0, 2, 4, 6, \dots, 2m-2)$ and note that C' is an m -cycle with $\ell(C') = \{2\}$.

For $0 \leq i \leq r-2$, let $P_i : 0, 9+8i, 4$ be the path of length 2 with edge lengths $9+8i, 5+8i$ and let $P'_i : 0, 11+8i, 4$ be the path of length 2 with edge lengths $11+8i, 7+8i$. Let $C_i = P_i \cup \rho^4(P_i) \cup \rho^8(P_i) \cup \dots \cup \rho^{2m-4}(P_i)$ and $C'_i = P'_i \cup \rho^4(P'_i) \cup \rho^8(P'_i) \cup \dots \cup \rho^{2m-4}(P'_i)$ and note that each is an m -cycle with $\ell(C_i) = \{5+8i, 9+8i\}$ and $\ell(C'_i) = \{7+8i, 11+8i\}$.

Finally, let $P'' : 0, 8r-3, -4$ be the path of length 2 with edge lengths $8r-3$ and $8r-1$. Let $C'' = P'' \cup \rho^4(P'') \cup \rho^8(P'') \cup \dots \cup \rho^{2m-4}(P'')$ and note that C'' is an m -cycle with $\ell(C'') = \{8r-3, 8r-1\}$.

Then $\{C, C', C_0, \dots, C_{r-2}, C'_0, \dots, C'_{r-2}, C''\}$ is a minimum generating set for a cyclic m -cycle system of $K_{2m} - I$. \square

We now consider the case when t is even and $t > 2$.

Lemma 3.2. *For each positive integer k and each positive integer $m \equiv 0 \pmod{8}$, there exists a cyclic m -cycle system of $K_{2mk} - I$.*

Proof. Let m and k be positive integers such that $m \equiv 0 \pmod{8}$. Lemma 3.1 handles the case when $k=1$ and thus we may assume that $k \geq 2$. Then $K_{2km} - I = \langle S' \rangle_{2km}$ where $S' = \{1, 2, \dots, km-1\}$. Since $K_{2m} - I$ has a cyclic m -cycle system by Lemma 3.1 and $\langle \{k, 2k, \dots, mk\} \rangle_{2km}$ consists of k vertex-disjoint copies of $K_{2m} - I$, we need only show that $\langle S \rangle_{2km}$ has a cyclic m -cycle system where $S = \{1, 2, \dots, mk\} \setminus \{k, 2k, \dots, mk\}$.

Let $A = [a_{i,j}]$ be the $(k-1) \times m$ array

$$\begin{bmatrix} k-1 & 2k-1 & 3k-1 & 4k-1 & & (m-1)k-1 & mk-1 \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 2 & k+2 & 2k+2 & 3k+2 & & (m-2)k+2 & (m-1)k+2 \\ 1 & k+1 & 2k+1 & 3k+1 & & (m-2)k+1 & (m-1)k+1 \end{bmatrix}.$$

It is straightforward to verify that A satisfies

$$\sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = \sum_{j \equiv 2,3 \pmod{4}} a_{i,j},$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}$$

for each i with $1 \leq i \leq k-1$.

For each $i = 1, 2, \dots, k-1$, the m -tuple

$(a_{i,1}, -a_{i,3}, a_{i,5}, -a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, -a_{i,6}, a_{i,4}, -a_{i,2}, a_{i,m})$ is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X = \{C_1, C_2, \dots, C_{k-1}\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S \rangle_{2km}$. \square

4. THE CASE WHEN $m \equiv 4 \pmod{8}$

In this section, we consider the case when $m \equiv 4 \pmod{8}$ and show that there exists a cyclic m -cycle system of $K_{mt} - I$ for each $t \geq 1$. We begin with the case when t is odd, say $t = 2k + 1$ for some nonnegative integer k .

Lemma 4.1. *For each nonnegative integer k and each $m \equiv 4 \pmod{8}$, there exists a cyclic m -cycle system of $K_{m(2k+1)} - I$.*

Proof. Let m and k be nonnegative integers such that $m \equiv 4 \pmod{8}$. Since $K_m - I$ has a cyclic hamiltonian cycle system [12], we may assume that $k \geq 1$. Let $m = 4r$ for some positive integer r . Then $K_{m(2k+1)} - I = \langle S' \rangle_{(2k+1)m}$ where $S' = \{1, 2, \dots, 4rk + 2r - 1\}$. Again, since $K_m - I$ has a cyclic hamiltonian cycle system [12] and $\langle \{2k+1, 4k+2, \dots, (2r-1)(2k+1)\} \rangle_{(2k+1)m}$ consists of $2k+1$ vertex-disjoint copies of $K_m - I$, we need only show that $\langle S \rangle_{(2k+1)m}$ has a cyclic m -cycle system where

$$S = \{1, 2, \dots, 4rk + 2r - 1\} \setminus \{2k+1, 4k+2, \dots, (2r-1)(2k+1)\}.$$

Let r and k be positive integers. Let $A = [a_{i,j}]$ be the $k \times m$ array

$$\begin{bmatrix} k & 2k & 3k+1 & 4k+1 & 5k+2 & & (4r-2)k+2r-2 & (4r-1)k+2r-1 & 4rk+2r-1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots & \vdots \\ 2 & k+2 & 2k+3 & 3k+3 & 4k+4 & & (4r-3)k+2r & (4r-2)k+2r+1 & (4r-1)k+2r+1 \\ 1 & k+1 & 2k+2 & 3k+2 & 4k+3 & & (4r-3)k+2r-1 & (4r-2)k+2r & (4r-1)k+2r \end{bmatrix}.$$

It is straightforward to verify that A satisfies

$$\sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = \sum_{j \equiv 2,3 \pmod{4}} a_{i,j},$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}$$

for each i with $1 \leq i \leq k$.

For each $i = 1, 2, \dots, k$, the m -tuple

$(a_{i,1}, -a_{i,3}, a_{i,5}, -a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, -a_{i,6}, a_{i,4}, -a_{i,2}, a_{i,m})$ is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X = \{C_1, C_2, \dots, C_k\}$ is a minimum generating set for a cyclic m -cycle system of $K_{m(2k+1)} - I$. \square

We now handle the case when t is even, say $t = 2k$ for some positive integer k .

Lemma 4.2. *For each positive integer k and each $m \equiv 4 \pmod{8}$, there exists a cyclic m -cycle system of $K_{2mk} - I$.*

Proof. As before, let m and k be positive integers such that $m \equiv 4 \pmod{8}$. Thus $m = 4r$ for some positive integer r . Then $K_{2mk} - I = \langle S' \rangle_{2km}$ where $S' = \{1, 2, \dots, 4rk - 1\}$. Since $K_m - I$ has a cyclic hamiltonian cycle system [12] and $\langle \{2k, 4k, \dots, (2r-1)(2k)\} \rangle_{2km}$ consists of $2k$ vertex-disjoint copies of $K_m - I$, we need only show that $\langle S \rangle_{2km}$ has a cyclic m -cycle system where

$$S = \{1, 2, \dots, 4rk - 1\} \setminus \{2k, 4k, \dots, (2r-1)(2k)\}.$$

Since $|S| = m(k-1) + m/2$, we will start by partitioning a subset $T \subseteq S$ with $|T| = m(k-1)$ into $k-1$ difference m -tuples.

Let $T = \{1, 2, \dots, 4rk - 1\} \setminus \{1, 2k, 4k - 1, 4k, 4k + 1, 6k, 8k - 1, 8k, 8k + 1, \dots, (4r-4)k - 1, (4r-4)k, (4r-4)k + 1, (4r-2)k, 4rk - 1\}$, and observe that $|T| = (k-1)m$. Let $A = [a_{i,j}]$, with entries from the set T , be the $(k-1) \times m$ array

$$\begin{bmatrix} k & 2k-1 & 3k-1 & 4k-2 & 5k & 6k-1 & 7k-1 & 8k-2 & 9k & & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \\ 3 & k+2 & 2k+2 & 3k+1 & 4k+3 & 5k+2 & 6k+2 & 7k+1 & 8k+3 & & \\ 2 & k+1 & 2k+1 & 3k & 4k+2 & 5k+1 & 6k+1 & 7k & 8k+2 & & \\ & & (4r-3)k & (4r-2)k-1 & (4r-1)k-1 & 4rk-2 & & & & & \\ \dots & \vdots & \vdots & \vdots & \vdots & \vdots & & & & & \\ & & (4r-4)k+3 & (4r-3)k+2 & (4r-2)k+2 & (4r-1)k+1 & & & & & \\ & & (4r-4)k+2 & (4r-3)k+1 & (4r-2)k+1 & (4r-1)k & & & & & \end{bmatrix}.$$

It is straightforward to verify that the array A satisfies

$$\sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = \sum_{j \equiv 2,3 \pmod{4}} a_{i,j},$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}$$

for each i with $1 \leq i \leq k-1$.

For each $i = 1, 2, \dots, k-1$, the m -tuple

$$(a_{i,1}, -a_{i,3}, a_{i,5}, -a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, -a_{i,6}, a_{i,4}, -a_{i,2}, a_{i,m})$$

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X = \{C_1, C_2, \dots, C_{k-1}\}$ is a minimum generating set for a cyclic m -cycle system of $\langle T \rangle_{2km}$.

It now remains to find a minimum generating set for a cyclic m -cycle system of $\langle B \rangle_{2km}$ where $B = \{1, 4k-1, 4k+1, 8k-1, 8k+1, \dots, (4r-4)k-1, (4r-4)k+1, 4rk-1\}$. For $i = 1, 2, \dots, r$, define $d_{2i-1} = 4(i-1)k+1$ and $d_{2i} = 4ik-1$. Observe that $B = \{d_1, d_2, \dots, d_{2r}\}$ and $d_{j+2} - d_j = 4k$ for $j = 1, 2, \dots, 2r-2$. Since $m \equiv 4 \pmod{8}$, it follows that r is odd. Let $P_1 : 0, 1, 4k$, and let $P_i : 0, d_{2i+1}, 4k$ if i is even and let $P_i : 0, d_{2i}, 4k$ if i is odd. Let $C'_i = P_i \cup \rho^{4k}(P_i) \cup \rho^{8k}(P_i) \cup \dots \cup \rho^{(2m-4)k}(P_i)$, and note that C'_i is an m -cycle with

$\ell(C'_1) = \{1, 4k - 1\}$, $\ell(C'_i) = \{d_{2i-1}, d_{2i+1}\}$ if i is even, and $\ell(C'_i) = \{d_{2i-2}, d_{2i}\}$ if i is odd. Then $\ell(C'_1) \cup \ell(C'_2) \cup \dots \cup \ell(C'_r) = B$ so that $\{C'_1, C'_2, \dots, C'_r\}$ is a minimum generating set for $\langle B \rangle_{2km}$. \square

5. THE CASE WHEN $m \equiv 2 \pmod{4}$

In this section, we consider the case when $m \equiv 2 \pmod{4}$ and prove parts (2) and (4) of Theorem 1.1. We divide this proof into three parts, each dealt with in its own subsection. First we consider the case $t \equiv 0 \pmod{4}$. Then we consider the case $m \equiv 2 \pmod{8}$ and $t \equiv 1 \pmod{4}$. Finally we consider the case $m \equiv 6 \pmod{8}$ and $t \equiv 3 \pmod{4}$.

5.1. The case when $t \equiv 0 \pmod{4}$.

We consider the case $t \equiv 0 \pmod{4}$, starting with the special case $t = 4$.

Lemma 5.1. *For each positive integer $m \geq 6$ with $m \equiv 2 \pmod{4}$, there exists a cyclic m -cycle system of $K_{4m} - I$.*

Proof. Let $m \geq 6$ be a positive integer with $m \equiv 2 \pmod{4}$. Then $K_{4m} - I = \langle S' \rangle_{4m}$ where $S' = \{1, 2, \dots, 2m - 1\}$. The proof proceeds as follows. We begin by finding one difference m -tuple which corresponds to an m -cycle C with $|\ell(C)| = m$. Note that $\langle \{4\} \rangle_{4m}$ consists of four vertex disjoint m -cycles. For the remaining $m - 2$ edge lengths in $S' \setminus (\ell(C) \cup \{4\})$, we find $(m - 2)/2$ paths P_i of length 2, ending at vertex 8 or -8 , so that $C_i = P_i \cup \rho^8(P_i) \cup \rho^{16}(P_i) \cup \dots \cup \rho^{4m-8}(P_i)$ is an m -cycle. Then this collection of cycles will give a minimum generating set for a cyclic m -cycle system of $K_{4m} - I$.

Consider the difference m -tuple

$$(1, -2, 6, -10, \dots, 2m - 6, -(2m - 2), -3, 8, -12, \dots, 2m - 12, -(2m - 8), 2m - 4)$$

and the corresponding m -cycle C with $\ell(C) = \{1, 2, 3, 6, 8, \dots, 2m - 2\}$. It is straightforward to verify that the odd vertices visited all lie between $-m + 1$ and $m - 1$ with no duplication. Similarly, the even vertices visited all lie between $-2m + 4$ and -4 , and have no duplication.

Let $C' = (0, 4, 8, \dots, 4m - 4)$ and note that C' is an m -cycle with $\ell(C') = \{4\}$.

Let $m = 8k + m'$, so m' is either 2 or 6. If $k = 0$, then $m' = 6$ and let $P : 0, 13, 8$ be the path of length 2 with edge lengths 11, 5. Then, $C'' = P \cup \rho^8(P) \cup \rho^{16}(P)$ is a 6-cycle with $\ell(C'') = \{11, 5\}$. Then $\{C, C', C''\}$ is a minimum generating set for cyclic 6-cycle system of $K_{24} - I$. Now suppose that $k \geq 1$. For $0 \leq i \leq k - 1$, let $P_i : 0, 13 + 16i, 8$ be the path of length 2 with edge lengths $13 + 16i, 5 + 16i$; let $P'_i : 0, 15 + 16i, 8$ be the path of length 2 with edge lengths $15 + 16i, 7 + 16i$; let $P''_i : 0, 17 + 16i, 8$ be the path of length 2 with edge lengths $17 + 16i, 9 + 16i$; and let $P'''_i : 0, 19 + 16i, 8$ with edge lengths $19 + 16i, 11 + 16i$. Let $C_i = P_i \cup \rho^8(P_i) \cup \rho^{16}(P_i) \cup \dots \cup \rho^{4m-8}(P_i)$, $C'_i = P'_i \cup \rho^8(P'_i) \cup \rho^{16}(P'_i) \cup \dots \cup \rho^{4m-8}(P'_i)$, $C''_i = P''_i \cup \rho^8(P''_i) \cup \rho^{16}(P''_i) \cup \dots \cup \rho^{4m-8}(P''_i)$, and $C'''_i = P'''_i \cup \rho^8(P'''_i) \cup \rho^{16}(P'''_i) \cup \dots \cup \rho^{4m-8}(P'''_i)$ and note that each is an m -cycle with $\ell(C_i) = \{5 + 16i, 13 + 16i\}$, $\ell(C'_i) = \{7 + 16i, 15 + 16i\}$, $\ell(C''_i) = \{9 + 16i, 17 + 16i\}$, and $\ell(C'''_i) = \{11 + 16i, 19 + 16i\}$.

If $m' = 2$, then $\{C, C', C_0, C'_0, C''_0, C'''_0, \dots, C_{k-1}, C'_{k-1}, C''_{k-1}, C'''_{k-1}\}$ is a minimum generating set for a cyclic m -cycle system of $K_{4m} - I$. If $m' = 6$, then let $P_k : 0, 2m - 1, -8$ and $P'_k : 0, 2m - 3, -8$ be paths of length 2 with $\ell(P_k) = \{2m - 1, 2m - 7\}$ and $\ell(P'_k) = \{2m - 3, 2m - 5\}$. Let $C_k = P_k \cup \rho^8(P_k) \cup \rho^{16}(P_k) \cup \dots \cup \rho^{4m-8}(P_k)$ and $C'_k = P'_k \cup \rho^8(P'_k) \cup \rho^{16}(P'_k) \cup \dots \cup \rho^{4m-8}(P'_k)$

and observe that each is an m -cycle with $\ell(C_k) = \{2m - 1, 2m - 7\}$ and $\ell(C'_k) = \{2m - 3, 2m - 5\}$. Thus, $\{C, C', C_0, C'_0, C''_0, C'''_0, \dots, C_{k-1}, C'_{k-1}, C''_{k-1}, C'''_{k-1}, C_k, C'_k\}$ is a minimum generating set for a cyclic m -cycle system of $K_{4m} - I$. \square

We now consider the case when $t \equiv 0 \pmod{4}$ with $t > 4$.

Lemma 5.2. *For each positive integer k and each positive integer $m \equiv 2 \pmod{4}$ with $m \geq 6$, there exists a cyclic m -cycle system of $K_{4mk} - I$.*

Proof. Let $m \geq 6$ and k be positive integers such that $m \equiv 2 \pmod{4}$. Lemma 5.1 handles the case when $k = 1$ and thus we may assume that $k \geq 2$. Then $K_{4km} - I = \langle S' \rangle_{4km}$ where $S' = \{1, 2, \dots, 2km - 1\}$. Since $K_{4m} - I$ has a cyclic m -cycle system by Lemma 5.1 and $\langle \{k, 2k, \dots, 2km\} \rangle_{4km}$ consists of k vertex-disjoint copies of $K_{4m} - I$, we need only show that $\langle S \rangle_{2km}$ has a cyclic m -cycle system where $S = \{1, 2, \dots, 2km\} \setminus \{k, 2k, \dots, 2km\}$.

Let $A = [a_{i,j}]$ be the $2k \times m$ array

$$\begin{bmatrix} 2k & 4k & 6k & 8k & & (m-1)2k & 2km \\ 2k-1 & 2k+1 & 6k-1 & 8k-1 & & (m-1)2k-1 & 2km-1 \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 2 & 4k-2 & 4k+2 & 6k+2 & & (m-2)2k+2 & (m-1)2k+2 \\ 1 & 4k-1 & 4k+1 & 6k+1 & & (m-2)2k+1 & (m-1)2k+1 \end{bmatrix}.$$

(Observe that the second column does not follow the same pattern as the others.)

Let A' be the $(2k-2) \times m$ array obtained from A by deleting rows 1 and $k+1$. Then the entries in A' are precisely the elements of S . Also, it is straightforward to verify that A' satisfies

$$a_{i,j} + a_{i,j+3} = a_{i,j+1} + a_{i,j+2}$$

for each positive integer $j \equiv 3 \pmod{4}$ with $j \leq m-3$,

$$a_{i,1} + a_{i,2} + a_{i,m-3} + a_{i,m-1} = a_{i,m-2} + a_{i,m},$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}$$

for each i with $1 \leq i \leq 2k-2$.

For each $i = 1, 2, \dots, 2k-2$, the m -tuple

$$(a_{i,1}, a_{i,2}, -a_{i,4}, a_{i,6}, -a_{i,8}, a_{i,10}, \dots, -a_{i,m-2}, -a_{i,m}, a_{i,m-3}, -a_{i,m-5}, a_{i,m-7}, \dots, a_{i,3}, a_{i,m-1})$$

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X = \{C_1, C_2, \dots, C_{2k-2}\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S \rangle_{4km}$. \square

What remains is to find cyclic m -cycle systems of $K_{mt} - I$ for the appropriate odd values of t , which we do in the following subsections.

5.2. The case when $m \equiv 2 \pmod{8}$ and $t \equiv 1 \pmod{4}$.

In this subsection, we find a cyclic m -cycle system of $K_{mt} - I$ when $m \equiv 2 \pmod{8}$ and $t \equiv 1 \pmod{4}$. We begin with two special cases, namely when $m = 10$ or $t = 5$.

Lemma 5.3. *For each positive integer $t \equiv 1 \pmod{4}$ with $t > 1$, there exists a cyclic 10-cycle system of $K_{10t} - I$.*

Proof. Let $t \equiv 1 \pmod{4}$ with $t > 1$, say $t = 4s + 1$ where $s \geq 1$. Then $K_{10t} - I = \langle S' \rangle_{10t}$ where $S' = \{1, 2, \dots, 20s + 4\}$. Consider the paths $P_1 : 0, 5t - 1, 2t$ and $P_2 : 0, 5t - 2, 2t$. Then, $\ell(P_1) = \{3t - 1, 5t - 1\}$ and $\ell(P_2) = \{3t - 2, 5t - 2\}$. For $i \in \{1, 2\}$, let $C_i = P_i \cup \rho^{2t}(P_i) \cup \rho^{4t}(P_i) \cup \dots \cup \rho^{8t}(P_i)$. Then clearly each C_i is an 10-cycle and $X = \{C_1, C_2\}$ is a minimum generating set for $\langle \{3t - 2, 3t - 1, 5t - 2, 5t - 1\} \rangle_{10t}$. Since $3t - 3 = 12s$ and $5t - 2 = 20s + 3$, it remains to find a cyclic 10-cycle system of $\langle S \rangle_{10t}$ where $S = \{1, 2, \dots, 12s, 12s + 3, 12s + 4, \dots, 20s + 2\}$. Let $A = [a_{i,j}]$ be the $2s \times 10$ array

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 8s+1 & 8s+3 & 12s+3 & 12s+4 & 12s+5 & 12s+6 \\ 5 & 6 & 7 & 8 & 8s+2 & 8s+4 & 12s+7 & 12s+8 & 12s+9 & 12s+10 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 8s-3 & 8s-2 & 8s-1 & 8s & 12s-2 & 12s & 20s-1 & 20s & 20s+1 & 20s+2 \end{bmatrix}$$

Clearly, for each i with $1 \leq i \leq 2s$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq 10)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,10}.$$

Thus the 10-tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,5}, a_{i,7}, -a_{i,9}, -a_{i,8}, a_{i,6}, -a_{i,4}, a_{i,10})$$

is a difference 10-tuple and corresponds to an 10-cycle C'_i with $\ell(C'_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,10}\}$. Hence, $X' = \{C'_1, C'_2, \dots, C'_{2s}\}$ is a minimum generating set for a cyclic 10-cycle system of $\langle S \rangle_{10t}$. \square

We now consider the case when $t = 5$.

Lemma 5.4. *For each positive integer $m \equiv 2 \pmod{8}$, there exists a cyclic m -cycle system of $K_{5m} - I$.*

Proof. Let m be a positive integer such that $m \equiv 2 \pmod{8}$, say $m = 8r + 2$ for some positive integer r . By Lemma 5.3, we may assume $r \geq 2$. Then $K_{5m} - I = \langle S' \rangle_{5m}$ where $S' = \{1, 2, \dots, 20r + 4\}$.

Let $2r = 6q + 4 + b$ for integers $q \geq 0$ and $b \in \{0, 2, 4\}$. Let a be a positive integer such that $1 + \log_2(q + 2) \leq a \leq 1 + \log_2(5q + 2)$, and note that a exists since if $q = 0$ then $\log_2(q + 2)$ is an integer, while if $q \geq 1$ then $2(q + 2) = 2q + 4 \leq 4q + 2 < 5q + 2$. For nonnegative integers i and j , define $d_{i,j} = 10(2r - i) + j$. Consider the path $P_{i,j} : 0, d_{i,j}, 5 \cdot 2^a$ and observe that $\ell(P_{i,j}) = \{10(2r - i) + j, 10(2r - i) + j - 5 \cdot 2^a\}$. If $0 < j < 10$, then $C_{i,j} = P_{i,j} \cup \rho^{10}(P_{i,j}) \cup \rho^{20}(P_{i,j}) \cup \dots \cup \rho^{5m-10}(P_{i,j})$ is an m -cycle since $m \equiv 2 \pmod{8}$ gives

$\gcd(5 \cdot 2^a, 5m) = 10$. Thus, if $0 < j < 10$, $\ell(C_{i,j}) = \{10(2r - i) + j, 10(2r - i) + j - 5 \cdot 2^a\}$.
Let

$$X = \{C_{0,j} \mid 1 \leq j \leq 4\} \cup \{C_{i,j} \mid 1 \leq i \leq q \text{ and } 1 \leq j \leq 6\} \cup \{C_{q+1,j} \mid 6 - b + 1 \leq j \leq 6\}$$

and let

$$\begin{aligned} B &= \{20r + j, 20r + j - 5 \cdot 2^a \mid 1 \leq j \leq 4\} \\ &\cup \{10(2r - i) + j, 10(2r - i) + j - 5 \cdot 2^a \mid 1 \leq i \leq q \text{ and } 1 \leq j \leq 6\} \\ &\cup \{10(2r - q - 1) + j, 10(2r - q - 1) + j - 5 \cdot 2^a \mid 6 - b + 1 \leq j \leq 6\}, \end{aligned}$$

where if $q = 0$ or $b = 0$, we take the corresponding sets to be empty as necessary. Now B will consist of $4r$ distinct lengths and X will be a minimum generating set for $\langle B \rangle_{5m}$ if $20r + 4 - 5 \cdot 2^a \leq 10(2r - q - 1) + 6 - b$. Note that $1 + \log_2(q + 2) \leq a \leq 1 + \log_2(5q + 2)$ gives $q + 2 \leq 2^{a-1} \leq 5q + 2$. So,

$$20r + 4 - [10(2r - q - 1) + 6 - b] = 10q + 8 + b \leq 10q + 12$$

and

$$(10q + 12)/10 < q + 2 \leq 2^{a-1}.$$

Thus $20r + 4 - 5 \cdot 2^a \leq 10(2r - q - 1) + 6 - b$ so that B consists of $4r$ distinct lengths, and X is a minimum generating set for $\langle B \rangle_{5m}$.

It remains to find a cyclic m -cycle system of $\langle S' \setminus B \rangle_{5m}$. The smallest length in B is $10(2r - q - 1) + 6 - b + 1 - 5 \cdot 2^a$, and we wish to show $10(2r - q - 1) + 6 - b - 5 \cdot 2^a \geq 12$. So,

$$10(2r - q - 1) + 6 - b - 12 = 20r - 10q - 16 - b \geq 20r - 10q - 20$$

and $(20r - 10q - 20)/10 \geq 2r - q - 2$. Now

$$2r - q - 2 = 5q + 2 + b \geq 5q + 2 \geq 2^{a-1}.$$

Hence, $10(2r - q - 1) + 6 - b - 5 \cdot 2^a \geq 12$. Since $|B| = 4r$, we have $|S' \setminus B| = 20r + 4 - 4r = 2(8r + 2)$. Now

$$\begin{aligned} S' \setminus B &= \{1, 2, \dots, 10(2r - q - 1) + 6 - b - 5 \cdot 2^a\} \cup \{10(2r - i) - 5 \cdot 2^a - 3, \\ &10(2r - i) - 5 \cdot 2^a - 2, 10(2r - i) - 5 \cdot 2^a - 1, 10(2r - i) - 5 \cdot 2^a \mid 0 \leq i \leq q\} \\ &\cup \{10(2r) + 5 - 5 \cdot 2^a, \dots, 10(2r - q - 1) + 6 - b\} \\ &\cup \{10(2r - i) - 3, 10(2r - i) - 2, 10(2r - i) - 1, 10(2r - i) \mid 0 \leq i \leq q\}. \end{aligned}$$

Note that each the sets $\{1, 2, \dots, 10(2r - q - 1) + 6 - b - 5 \cdot 2^a\}$, $\{10(2r - i) - 5 \cdot 2^a - 3, 10(2r - i) - 5 \cdot 2^a - 2, 10(2r - i) - 5 \cdot 2^a - 1, 10(2r - i) - 5 \cdot 2^a \mid 0 \leq i \leq q\}$, $\{10(2r) + 5 - 5 \cdot 2^a, \dots, 10(2r - q - 1) + 6 - b\}$, and $\{10(2r - i) - 3, 10(2r - i) - 2, 10(2r - i) - 1, 10(2r - i) \mid 0 \leq i \leq q\}$ has even cardinality and consists of consecutive integers. Therefore, we may partition $S' \setminus B$ into sets $T, S_1, S_2, \dots, S_{8r-4}$ where $T = \{1, 2, \dots, 12\}$ and for $i = 1, 2, \dots, 8r - 4$, let $S_i = \{b_i, b_i + 1\}$ with $b_1 < b_2 < \dots < b_{8r-4}$.

Let $A = [a_{i,j}]$ be the $2 \times m$ array

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 9 & 11 & b_1 & b_1 + 1 & b_2 & b_2 + 1 & \cdots & b_{4r-2} & b_{4r-2} + 1 \\ 5 & 6 & 7 & 8 & 10 & 12 & b_{4r-1} & b_{4r-1} + 1 & b_{4r} & b_{4r} + 1 & \cdots & b_{8r-4} & b_{8r-4} + 1 \end{bmatrix}$$

It is straightforward to verify that, for $1 \leq i \leq 2$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq m)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}.$$

Hence, for $1 \leq i \leq 2$, the m -tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,5}, a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, a_{i,6}, -a_{i,4}, a_{i,m})$$

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X' = \{C_1, C_2\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S' \setminus B \rangle_{5m}$. \square

We are now ready to prove the main result of this subsection, namely, that $K_{mt} - I$ has a cyclic m -cycle system for every $t \equiv 1 \pmod{4}$ and $m \equiv 2 \pmod{8}$ with $t > 1$ if $m = 2p^\alpha$ for some prime p and integer $\alpha \geq 1$.

Lemma 5.5. *For each positive integer $t \equiv 1 \pmod{4}$ and each $m \equiv 2 \pmod{8}$ with $t > 1$ if $m = 2p^\alpha$ for some prime p and integer $\alpha \geq 1$, there exists a cyclic m -cycle system of $K_{mt} - I$.*

Proof. Let m and t be positive integers such that $m \equiv 2 \pmod{8}$ and $t \equiv 1 \pmod{4}$. Thus $m = 8r + 2$ for some positive integer r . Then $K_{mt} - I = \langle S' \rangle_{mt}$ where $S' = \{1, 2, \dots, (mt - 2)/2\}$. Since $K_m - I$ has a cyclic hamiltonian cycle system [12] if and only if $m \neq 2p^\alpha$ for some prime p and integer $\alpha \geq 1$, we may assume that $t > 1$. Thus, let $t = 4s + 1$ for some positive integer s . By Lemmas 5.3 and 5.4, we may assume that $s \geq 2$ and $r \geq 2$.

The proof proceeds as follows. We begin by finding a set $B \subseteq S'$ such that $|B| = 4r$ and $\langle B \rangle_{mt}$ has a cyclic m -cycle system with a minimum generating set X consisting of cycles each with two distinct lengths and orbit $2t$. We then construct an $(|S' \setminus B|/m) \times m$ array $A = [a_{i,j}]$ with the property that for each i with $1 \leq i \leq |S' \setminus B|/m$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq m)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}.$$

Thus for each $i = 1, 2, \dots, |S' \setminus B|/m$, the m -tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,5}, a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, a_{i,6}, -a_{i,4}, a_{i,m})$$

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X' = \{C_1, C_2, \dots, C_{|S' \setminus B|/m}\}$ will be a minimum generating set for a cyclic m -cycle system of $\langle S' \setminus B \rangle_{mt}$.

Let $w = \lfloor r/2 \rfloor$, and let $\delta_r = 2(r/2 - w)$, so that $\delta_r = 1$ if r is odd and $\delta_r = 0$ if r is even. Write $w = qs + b$ where q and b are non-negative integers with $0 \leq b < s$ (note that it may be the case that $q = 0$). For integers i and j , define $d_{i,j} = 4(r - 2i)t + j$. Consider the path $P_{i,j} : 0, d_{i,j}, 4t$ and observe that $\ell(P_{i,j}) = \{4(r - 2i)t + j, 4(r - 2i - 1)t + j\}$. If $0 < j < t$,

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X' = \{C_1, C_2, \dots, C_{2s}\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S' \setminus B \rangle_{mt}$. \square

5.3. The Case when $m \equiv 6 \pmod{8}$ and $t \equiv 3 \pmod{4}$.

In this subsection, we find a cyclic m -cycle system of $K_{mt} - I$ when $m \equiv 6 \pmod{8}$ and $t \equiv 3 \pmod{4}$. We begin with three special cases, namely when $m = 6$, $m = 14$, or $t = 3$. We first consider the case $m = 6$.

Lemma 5.6. *For all positive integers $t \equiv 3 \pmod{4}$, there exists a cyclic 6-cycle system of $K_{6t} - I$.*

Proof. Let t be a positive integer such that $t \equiv 3 \pmod{4}$, say $t = 4s + 3$ for some non-negative integer s . Then $K_{6t} - I = \langle S' \rangle_{6t}$ where $S' = \{1, 2, \dots, 12s + 8\}$.

Consider the paths $P_i : 0, 3t - i, 2t$, for $1 \leq i \leq 4$; then $\ell(P_i) = \{3t - i, t - i\}$. Next, let $C_i = P_i \cup \rho^{2t}(P_i) \cup \rho^{4t}(P_i)$. Then each C_i is a 6-cycle and $X = \{C_1, C_2, C_3, C_4\}$ is a minimum generating set for $\langle B \rangle_{6t}$ where $B = \{3t - i, t - i \mid 1 \leq i \leq 4\}$. Now, $t - 5 = 4s - 2$ and thus $S' \setminus B = \{1, 2, \dots, 4s - 2, 4s + 3, 4s + 4, \dots, 12s + 4\}$, and so we must find a cyclic 6-cycle system of $\langle S' \setminus B \rangle_{6t}$. Let $A = [a_{i,j}]$ be the $2s \times 6$ array

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 8s+5 & 8s+7 \\ 5 & 6 & 7 & 8 & 8s+6 & 8s+8 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 4s-3 & 4s-2 & 4s+3 & 4s+4 & \alpha & \alpha+2 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 8s+1 & 8s+2 & 8s+3 & 8s+4 & 12s+2 & 12s+4 \end{bmatrix}$$

where

$$\alpha = \begin{cases} 10s + 2 & \text{if } s \text{ is even,} \\ 10s + 3 & \text{if } s \text{ is odd.} \end{cases}$$

Clearly, for each i with $1 \leq i \leq 2s$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq 6)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,6}.$$

Thus the 6-tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,4}, -a_{i,5}, a_{i,6})$$

is a difference 6-tuple and corresponds to a 6-cycle C'_i with $\ell(C'_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,6}\}$. Hence, $X' = \{C'_1, C'_2, \dots, C'_{2s}\}$ is a minimum generating set for a cyclic 6-cycle system of $\langle S' \setminus B \rangle_{6t}$. \square

Next we consider the case when $m = 14$.

Lemma 5.7. *For all positive integers $t \equiv 3 \pmod{4}$, there exists a cyclic 14-cycle system of $K_{14t} - I$.*

Proof. Let t be a positive integer such that $t \equiv 3 \pmod{4}$, say $t = 4s+3$ for some non-negative integer s . Then $K_{14t} - I = \langle S' \rangle_{14t}$ where $S' = \{1, 2, \dots, 28s + 20\}$.

Consider the paths $P_i : 0, 7t - i, 2t$, for $1 \leq i \leq 10$; then $\ell(P_i) = \{7t - i, 5t - i\}$. Next, let $C_i = P_i \cup \rho^{2t}(P_i) \cup \rho^{4t}(P_i) \cup \dots \cup \rho^{12t}(P_i)$. Then each C_i is a 14-cycle and $X = \{C_1, C_2, \dots, C_{10}\}$ is a minimum generating set for $\langle B \rangle_{14t}$ where $B = \{7t - i, 5t - i \mid 1 \leq i \leq 10\}$. Now, $5t - 10 = 20s + 5$ and thus $S' \setminus B = \{1, 2, \dots, 20s + 4, 20s + 15, 20s + 16, \dots, 28s + 10\}$, and so we must find a cyclic 14-cycle system of $\langle S' \setminus B \rangle_{14t}$. Let $A = [a_{i,j}]$ be the $2s \times 14$ array

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 8s+1 & 8s+3 & 12s+1 & 12s+2 & 12s+3 & 12s+4 \\ 5 & 6 & 7 & 8 & 8s+2 & 8s+4 & 12s+5 & 12s+6 & 12s+7 & 12s+8 \\ 9 & 10 & 11 & 12 & 8s+5 & 8s+7 & 12s+9 & 12s+10 & 12s+11 & 12s+12 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 8s-3 & 8s-2 & 8s-1 & 8s & 12s-2 & 12s & 20s-3 & 20s-2 & 20s-1 & 20s \end{bmatrix} \cdot \begin{bmatrix} 20s+1 & 20s+2 & 20s+3 & 20s+4 \\ 20s+15 & 20s+16 & 20s+17 & 20s+18 \\ 20s+19 & 20s+20 & 20s+21 & 20s+22 \\ \vdots & \vdots & \vdots & \vdots \\ 28s+7 & 28s+8 & 28s+9 & 28s+10 \end{bmatrix}.$$

Clearly, for each i with $1 \leq i \leq 2s$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq 14)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,14}.$$

Thus the 14-tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,5}, a_{i,7}, -a_{i,9}, a_{i,11}, -a_{i,13}, -a_{i,12}, a_{i,10}, -a_{i,8}, a_{i,6}, -a_{i,4}, a_{i,14})$$

is a difference 14-tuple and corresponds to a 14-cycle C'_i with $\ell(C'_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,14}\}$. Hence, $X' = \{C'_1, C'_2, \dots, C'_{2s}\}$ is a minimum generating set for a cyclic 14-cycle system of $\langle S' \setminus B \rangle_{14t}$. \square

We now consider the case when $t = 3$.

Lemma 5.8. *For all positive integers $m \equiv 6 \pmod{8}$, there exists a cyclic m -cycle system of $K_{3m} - I$.*

Proof. Let m be a positive integer such that $m \equiv 6 \pmod{8}$, say $m = 8r + 6$ for some non-negative integer r . By Lemmas 5.6 and 5.7, we may assume $r \geq 2$. Then $K_{3m} - I = \langle S' \rangle_{mt}$ where $S' = \{1, 2, \dots, 12r + 8\}$. Write $2r = 4q + b + 2$ for integers $q \geq 0$ and $b \in \{0, 2\}$, and let a be a positive integer such that $1 + \log_2(q + 1) \leq a \leq 1 + \log_2(3q + 4/3 + 5b/6)$. For integers i and j , define $d_{i,j} = 6(2r - i) + j$. Then consider the path $P_{i,j} : 0, d_{i,j}, 3 \cdot 2^a$; so $\ell(P_{i,j}) = \{6(2r - i) + j, 6(2r - i) + j - 3 \cdot 2^a\}$. Now, let $C_{i,j} = P_{i,j} \cup \rho^6(P_{i,j}) \cup \dots \cup \rho^{3(m-2)}(P_{i,j})$. Then $C_{i,j}$ is an m -cycle since $m \equiv 6 \pmod{8}$ implies $\gcd(3 \cdot 2^a, 3m) = 6$. Thus, $\ell(C_{i,j}) = \ell(P_{i,j})$.

Now, let

$$\begin{aligned} X &= \{C_{0,j} \mid j = 7, 8\} \\ &\cup \{C_{i,j} \mid 0 \leq i \leq q-1 \text{ and } 1 \leq j \leq 4\} \\ &\cup \{C_{q,j} \mid 5-b \leq j \leq 4\} \end{aligned}$$

and let

$$\begin{aligned} B &= \{12r+7, 12r+7-3 \cdot 2^a, 12r+8, 12r+8-3 \cdot 2^a\} \\ &\cup \{6(2r-i)+j, 6(2r-i)-3 \cdot 2^a+j \mid 0 \leq i \leq q-1 \text{ and } 1 \leq j \leq 4\} \\ &\cup \{6(2r-q)+j, 6(2r-q)-3 \cdot 2^a+j \mid 5-b \leq j \leq 4\} \end{aligned}$$

where, if $q = 0$ or $b = 0$, we take the corresponding sets to be empty as necessary. Now B will consist of $4r$ distinct lengths and X will be a minimum generating set for $\langle B \rangle_{3m}$ if $12r+8-3 \cdot 2^a \leq 6(2r-q)+5-b-1$. Note that $1 + \log_2(q+1) \leq a$ so that $q+1 \leq 2^{a-1}$. Next,

$$12r+8 - [6(2r-q) + 5 - b - 1] = 6q + 4 + b \leq 6q + 6 = 6(q+1) \leq 6 \cdot 2^{a-1} = 3 \cdot 2^a,$$

and hence $12r+8-3 \cdot 2^a \leq 6(2r-q)+5-b-1$. Thus, B consists of $4r$ distinct lengths, and X is a minimum generating set for $\langle B \rangle_{3m}$. Now, the smallest length in B is $6(2r-q)+5-b-3 \cdot 2^a$ and we want this length to be greater than 8. Recall that $a \leq 1 + \log_2(3q+3/2+5b/6)$ and thus $2^{a-1} \leq 3q+3/2+5b/6$. Hence, $3 \cdot 2^a \leq 18q+9+5b = 12r-6q-3-b$ since $2r = 4q+b+2$. Therefore, $6(2r-q)+5-b-3 \cdot 2^a \geq 8$. Since $|B| = 4r$, we have $|S' \setminus B| = 8r+8$. Note that

$$\begin{aligned} S' \setminus B &= \{1, 2, \dots, 6(2r-q) + 5 - b - 3 \cdot 2^a - 1\} \\ &\cup \{6(2r-i) - 3 \cdot 2^a + 5, 6(2r-i) - 3 \cdot 2^a + 6 \mid 0 \leq i \leq q\} \\ &\cup \{12r - 3 \cdot 2^a + 9, \dots, 6(2r-q) + 5 - b - 1\} \\ &\cup \{6(2r-i) + 5, 6(2r-i) + 6 \mid 0 \leq i \leq q\}. \end{aligned}$$

Note that $S' \setminus B$ has been written as the disjoint union of sets, each of which has even cardinality and consists of consecutive integers. Therefore, we may partition $S' \setminus B$ into sets $T, S_1, S_2, \dots, S_{4r}$ where $T = \{1, 2, \dots, 8\}$ and for $i = 1, 2, \dots, 4r$, let $S_i = \{b_i, b_i + 1\}$ with $b_1 < b_2 < \dots < b_{4r}$.

Consider the m -tuple

$$\begin{aligned} (1, -3, 6, -7, b_1, -b_2, b_3, -b_4, \dots, b_{4r-1}, -b_{4r}, -(b_{4r-1} + 1), b_{4r-2} + 1, \\ -(b_{4r-3} + 1), b_{4r-4} + 1, \dots, b_2 + 1, -(b_1 + 1), 8, -5, b_{4r} + 1) \end{aligned}$$

which is a difference m -tuple and corresponds to an m -cycle C_1 with

$$\ell(C_1) = \{1, 3, 5, 6, 7, 8, b_1, b_1 + 1, b_2, b_2 + 1, \dots, b_{4r}, b_{4r} + 1\}.$$

Then consider the path $P : 0, 2, 6$; so $\ell(P) = \{2, 4\}$. Now, let $C_2 = P \cup \rho^6(P) \cup \dots \cup \rho^{3(m-2)}(P)$. Then C_2 is an m -cycle since $m \equiv 6 \pmod{8}$ implies $\gcd(6, 3m) = 6$. Thus, $\ell(C_2) = \ell(P) = \{2, 4\}$. Hence, $X' = \{C_1, C_2\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S' \setminus B \rangle_{3m}$. \square

We now prove the main result of this subsection, namely that $K_{mt} - I$ has a cyclic m -cycle system for every $t \equiv 3 \pmod{4}$ and $m \equiv 6 \pmod{8}$.

Lemma 5.9. *For all positive integers $t \equiv 3 \pmod{4}$ and $m \equiv 6 \pmod{8}$, there exists a cyclic m -cycle system of $K_{mt} - I$.*

Proof. Let m and t be positive integers such that $m \equiv 6 \pmod{8}$ and $t \equiv 3 \pmod{4}$. Then $m = 8r + 6$ and $t = 4s + 3$ for some non-negative integers r and s . Then $K_{mt} - I = \langle S' \rangle_{mt}$ where $S' = \{1, 2, \dots, (4r + 3)t - 1\}$.

By Lemmas 5.6, 5.7, and 5.8, we may assume $s \geq 1$ and $r \geq 2$. First, write $6r + 4 = (2t - 2)q + (t - 1)\ell + b$ for integers q, ℓ and b with $q \geq 0, 0 \leq b < 2t - 2$, and $\ell = 0$ if $6r + 4 < t - 1$, or $\ell = 1$ otherwise. For integers i and j , define $d_{i,j} = 2t(2r - 2i - 1) + j$. Consider the path $P_{i,j} : 0, d_{i,j}, 2t$ and note that $\ell(P_{i,j}) = \{2t(2r - 2i - 1) + j, 2t(2r - 2i - 2) + j\}$. If $0 < j < 2t$, then $C_{i,j} = P_{i,j} \cup \rho^{2t}(P_{i,j}) \cup \rho^{4t}(P_{i,j}) \cup \dots \cup \rho^{(m-2)t}(P_{i,j})$ is an m -cycle since $m \equiv 6 \pmod{8}$ implies $\gcd(2t, mt) = 2t$. Thus, if $0 < j < 2t$, then $\ell(C_{i,j}) = \ell(P_{i,j})$.

Now, let

$$\begin{aligned} X &= \{C_{-1,j} \mid 1 \leq j \leq t - 1\} \\ &\cup \{C_{i,j} \mid 0 \leq i \leq q - 1 \text{ and } 1 \leq j \leq 2t - 2\} \\ &\cup \{C_{q,j} \mid 2t - 1 - b \leq j \leq 2t - 2\} \end{aligned}$$

and let

$$\begin{aligned} B &= \{2t(2r + 1) + j, 2t(2r) + j \mid 1 \leq j \leq t - 1\} \\ &\cup \{2t(2r - 2i - 1) + j, 2t(2r - 2i - 2) + j \mid 1 \leq j \leq 2t - 2 \text{ and } 0 \leq i \leq q - 1\} \\ &\cup \{2t(2r - (2q + 1)) + 2t - 1 - b + j, 2t(2r - 2q - 2) + 2t - 1 - b + j \mid 0 \leq j \leq b - 1\}. \end{aligned}$$

where we take the first set to be empty if $\ell = 0$, the second to be empty if $q = 0$, and the third to be empty if $b = 0$. Then X is a minimum generating set for $\langle B \rangle_{mt}$.

Now we must find a cyclic m -cycle system of $\langle S' \setminus B \rangle_{mt}$. First, $|B| = 2[(2t - 2)q + (t - 1)\ell + b] = 12r + 8$ so that $|S' \setminus B| = (4r + 3)t - 1 - 12r - 8 = (8r + 6)(2s)$. Moreover,

$$\begin{aligned} S' \setminus B &= \{1, 2, \dots, 2t(2r - 2q - 1) - b - 2\} \\ &\cup \{2t(2r - 2q - 1) - 1, 2t(2r - 2q - 1), \dots, 2t(2r - 2q) - b - 2\} \\ &\cup \{2t(2r - i) - 1, 2t(2r - i) \mid 0 \leq i \leq 2q\} \\ &\cup \{4rt + t, 4rt + t + 1, \dots, 4rt + 2t\}. \end{aligned}$$

The smallest length in B is $4t(r - q - 1) + (2t - 1) - b$, and we must verify that this length is at least $12s + 1$. Note that we have $2t - 1 - b > 1$. Thus, it is sufficient to prove that $4t(r - q - 1) \geq 12s$, or $t(r - q - 1) \geq 3s$. This inequality follows if $r > q + 1$. Clearly, this is true if $q = 0$ since $r \geq 2$, so assume $q \geq 1$. Then $\ell = 1$, and so $6r + 4 = 2q(4s + 2) + (4s + 2) + b$,

or

$$\begin{aligned} 3r + 2 &= q(4s + 2) + 2s + 1 + b/2 \\ &= 4qs + 2q + 2s + 1 + b/2 \\ &\geq 6q + 3 \quad (\text{since } s \geq 1). \end{aligned}$$

So, $r \geq 2q + 1/3 > q + 1$ since $q \geq 1$. Since the smallest length in B is at least $12s + 1$ and $S' \setminus B$ consists of sets of consecutive integers of even cardinality, we may partition $S' \setminus B$ into sets T, S_1, \dots, S_{8rs} where $T = \{1, 2, \dots, 12s\}$, and for $i = 1, 2, \dots, 8rs$, $S_i = \{b_i, b_i + 1\}$ with $b_1 \leq b_2 \leq \dots \leq b_{8rs}$. Let $A = [a_{i,j}]$ be the $2s \times m$ array

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 8s+1 & 8s+3 & b_1 & b_1+1 \\ 5 & 6 & 7 & 8 & 8s+2 & 8s+4 & b_{4r+1} & b_{4r+1}+1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \dots \\ 8s-3 & 8s-2 & 8s-1 & 8s & 12s-2 & 12s & b_{8rs-4r+1} & b_{8rs-4r+1}+1 \\ & & & & & & & & \\ & b_2 & b_2+1 & \dots & b_{4r} & b_{4r}+1 \\ & b_{4r+2} & b_{4r+2}+1 & \dots & b_{8r} & b_{8r}+1 \\ & \vdots & \vdots & \dots & \vdots & \vdots \\ & b_{8rs-4r+2} & b_{8rs-4r+2}+1 & \dots & b_{8rs} & b_{8rs}+1 \end{bmatrix}.$$

Clearly, for each i with $1 \leq i \leq 2s$,

$$a_{i,2} + \sum_{j \equiv 0,1 \pmod{4}} a_{i,j} = a_{i,1} + \sum_{j \equiv 2,3 \pmod{4}} a_{i,j} \quad (\text{where } 3 \leq j \leq m)$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,m}.$$

Thus the m -tuple

$$(a_{i,1}, -a_{i,2}, a_{i,3}, -a_{i,5}, a_{i,7}, \dots, a_{i,m-3}, -a_{i,m-1}, -a_{i,m-2}, a_{i,m-4}, -a_{i,m-6}, \dots, a_{i,6}, -a_{i,4}, a_{i,m})$$

is a difference m -tuple and corresponds to an m -cycle C_i with $\ell(C_i) = \{a_{i,1}, a_{i,2}, \dots, a_{i,m}\}$. Hence, $X' = \{C_1, C_2, \dots, C_{2s}\}$ is a minimum generating set for a cyclic m -cycle system of $\langle S' \setminus B \rangle_{mt}$. \square

REFERENCES

- [1] B. Alspach and H. Gavlas, Cycle decompositions of K_n and $K_n - I$, *J. Combin. Theory Ser. B*, **81**(2001), 77–99.
- [2] A. Blinco, S. El-Zanati and C. Vanden Eynden, On the cyclic decomposition of complete graphs into almost-bipartite graphs, *Discrete Math.* **284** (2004), 71–81.
- [3] D. Bryant, H. Gavlas and A. Ling, Skolem-type difference sets for cycles, *Electronic J. Combin.* **10** (2003), #R38.
- [4] D. Bryant and C. A. Rodger, Cycle Decompositions, in the *CRC Handbook of Combinatorial Designs, 2nd Ed.*, (eds. C.J. Colbourn and J.H. Dinitz), Chapman & Hall/CRC, Boca Raton FL (2007).
- [5] M. Buratti, Cycle decompositions with a sharply vertex transitive automorphism group, *Le Matematiche* **59** (2004), 91–105.
- [6] M. Buratti and A. Del Fra, Existence on cyclic k -cycle systems of the complete graph. *Discrete Math.* **261** (2003), 113–125.

- [7] M. Buratti and A. Del Fra, Cyclic Hamiltonian cycle systems of the complete graph, *Discrete Math.*, **279** (2004), 107–119.
- [8] M. Buratti and G. Rinaldi, A non-existence result on cyclic cycle-decompositions of the cocktail party graph, *Discrete Math.* **309** (2009) 4722–4726.
- [9] J. D. Dixon and B. Mortimer, *Permutation Groups*, Springer-Verlag New York, Berlin, Heidelberg, Graduate Texts in Mathematics, **163**, 1996.
- [10] S.I. El-Zanati, N. Punnim, C. Vanden Eynden, On the cyclic decomposition of complete graphs into bipartite graphs, *Austral. J. Combin.* **24** (2001), 209–219.
- [11] H.-L. Fu and S.-L. Wu, Cyclically decomposing the complete graph into cycles, *Discrete Math.* **282** (2004), 267–273.
- [12] H. Jordon and J. Morris, Cyclic hamiltonian cycle systems of the complete graph minus a 1-factor, *Discrete Math.* **308** (2008), 2440–2449.
- [13] A. Kotzig, On decompositions of the complete graph into $4k$ -gons, *Mat.-Fyz. Cas.* **15** (1965), 227–233.
- [14] R. Peltesohn, Eine Lösung der beiden Heffterschen Differenzenprobleme, *Compos. Math.* **6** (1938), 251–257.
- [15] A. Rosa, On cyclic decompositions of the complete graph into $(4m+2)$ -gons, *Mat.-Fyz. Cas.* **16** (1966), 349–352.
- [16] A. Rosa, On the cyclic decompositions of the complete graph into polygons with an odd number of edges, *Časopis Pěst. Math.* **91** (1966), 53–63.
- [17] M. Šajna, Cycle Decompositions III: Complete graphs and fixed length cycles, *J. Combin. Des.* **10** (2002), 27–78.
- [18] A. Veitri, Cyclic k -cycles systems of order $2kn+k$; a solution of the last open cases, *J. Combin. Des.* **12** (2004), 299–310.
- [19] S.-L. Wu, Cyclic even cycle systems of the complete graph, *J. Combin. Des.* **20** (2012) 23–29.
- [20] S.-L. Wu and H.-L. Fu, Cyclic m -cycle systems with $m \leq 32$ or $m = 2q$ with q a prime power, *J. Combin. Des.* **14** (2006), 66–81.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ALBION COLLEGE, ALBION, MI 49224, USA

E-mail address: hjordon@albion.edu

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, UNIVERSITY OF LETHBRIDGE, LETHBRIDGE, AB, CANADA T1K 3M4

E-mail address: joy@cs.uleth.ca