

# SIMULTANEOUS NON-VANISHING OF TWISTS

AMIR AKBARY

ABSTRACT. Let  $f$  be a newform of even weight  $k$ , level  $M$  and character  $\psi$  and let  $g$  be a newform of even weight  $l$ , level  $N$  and character  $\eta$ . We give a generalization of a theorem of Elliott, regarding the average values of Dirichlet  $L$ -functions, in the context of twisted modular  $L$ -functions associated to  $f$  and  $g$ . Using this result, we find a lower bound in terms of  $Q$  for the number of primitive Dirichlet characters modulo prime  $q \leq Q$  whose twisted product  $L$ -functions  $L_{f,\chi}(s)L_{g,\chi}(s)$  are non-vanishing at a fixed point  $s_0 = \sigma_0 + it_0$  with  $\frac{1}{2} < \sigma_0 \leq 1$ .

## 1. INTRODUCTION

Let  $L_\chi(s) = \sum_{n=1}^{\infty} \chi(n)n^{-s}$  be the Dirichlet  $L$ -function associated to a Dirichlet character  $\chi$ . In [E], Elliott proved the following.

**Theorem** *Let  $Q \geq 2$  be a real number, and  $s_0 = \sigma_0 + it_0$  a complex number in the half-plane  $\sigma_0 > \frac{1}{2}$ . Then we have*

$$\sum_{p \leq Q} \sum_{\chi \neq \chi_0} |L_\chi(s_0)|^2 = \frac{Q^2}{2 \log Q} \zeta(2\sigma_0) + O\left(\frac{Q^2}{(\log Q)^2}\right)$$

as  $Q \rightarrow \infty$ . Here the inner sum is taken over all non-principal characters (mod  $p$ ), for each prime  $p$ , and the outer sum over all prime numbers not exceeding  $Q$ .

Our first goal in this paper is to give a generalization of this theorem in the context of twisted modular  $L$ -functions. Let  $S_k(\Gamma_0(M), \psi)$  be the space of holomorphic cusp forms of even weight  $k$ , level  $M$  and character  $\psi$ . For  $f \in S_k(\Gamma_0(M), \psi)$ , let

$$f(z) = \sum_{n=1}^{\infty} a_f(n) n^{\frac{k-1}{2}} e^{2\pi i n z}$$

be the Fourier expansion of  $f$  at  $i\infty$ . Let  $\chi$  be a primitive Dirichlet character mod  $q$  with  $(q, M) = 1$ . Then the twisted  $L$ -function associated to  $f$  and  $\chi$  is defined (for  $\text{Re}(s) > 1$ ) by

$$L_{f,\chi}(s) = \sum_{n=1}^{\infty} \frac{a_f(n)\chi(n)}{n^s}.$$

Let

$$L_{\infty,k}(s) = (2\pi)^{-s} \Gamma\left(\frac{k-1}{2} + s\right),$$

and

$$\Lambda_{f,\chi}(s) = (q\sqrt{M})^s L_{\infty,k}(s) L_{f,\chi}(s).$$

Then it is known that  $\Lambda_{f,\chi}(s)$  is entire and if  $f$  is a newform (in Atkin-Lehner sense) it satisfies the functional equation

---

2000 *Mathematics Subject Classification.* 11F67.

Research partially supported by NSERC.

$$(1) \quad \Lambda_{f,\chi}(s) = \epsilon_{f,\chi} \Lambda_{\bar{f},\bar{\chi}}(1-s)$$

where  $\bar{f}$  is the conjugate newform in  $S_k(\Gamma_0(M), \bar{\psi})$ . Here  $\epsilon_{f,\chi} = \epsilon_f \psi(q) \chi(M) \tau(\chi)^2 q^{-1}$  where  $|\epsilon_f| = 1$  and  $\tau(\chi)$  is the Gauss sum. Note that  $|\epsilon_{f,\chi}| = 1$ .

We recall that for  $f \in S_k(\Gamma_0(M), \psi)$  and  $g \in S_l(\Gamma_0(N), \eta)$  the Rankin-Selberg convolution  $L$ -function is defined (for  $\operatorname{Re}(s) > 1$ ) by

$$L(f \otimes g, s) = \sum_{n=1}^{\infty} \frac{a_f(n) \bar{b}_g(n)}{n^s}.$$

The following can be considered as a modular analogue of the above theorem of Elliott.

**Theorem 1.1.** *Let  $f \in S_k(\Gamma_0(M), \psi)$  and  $g \in S_l(\Gamma_0(N), \eta)$  be newforms. Let  $Q \geq 2$  and  $s_0 = \sigma_0 + it_0$  be a complex number with  $\sigma_0 > \frac{1}{2}$ . Then we have*

$$\sum_{\substack{q \leq Q, q \text{ prime} \\ (q, MN)=1}} \sum_{\chi \pmod{q}}^* L_{f,\chi}(s_0) \overline{L_{g,\chi}(s_0)} = \frac{Q^2}{2 \log Q} \frac{\phi(MN)}{MN} L(f \otimes g, 2\sigma_0) + O\left(\frac{Q^2}{(\log Q)^2}\right)$$

where the inner sum is taken over the primitive characters modulo prime  $q$ . The implied constant depends on  $f, g$  and  $s_0$ . Here,  $\phi$  is the Euler function.

In proving Theorem 1.1, we first find an asymptotic formula for the values  $L_{f,\chi}(s_0) \overline{L_{g,\chi}(s_0)}$  on average when  $\chi$  varies on the set of primitive characters modulo fixed positive integer  $q$  (see Proposition 2.5). This result generalizes a theorem of Stefanicki ([S], Theorem 2(a)).

Theorem 1.1 has an interesting application in the problem of non-vanishing of twisted  $L$ -functions inside the critical strip. In Proposition 3.1, by employing the large sieve inequality for characters, we establish an upper bound for the mean square of the values  $|L_{f,\chi}(s_0) \overline{L_{g,\chi}(s_0)}|$ . Together, Theorem 1.1 and Proposition 3.1 imply the following.

**Theorem 1.2.** *Let  $f \in S_k(\Gamma_0(M), \psi)$  and  $g \in S_l(\Gamma_0(N), \eta)$  be newforms. Let  $s_0 = \sigma_0 + it_0$  be a fixed point in the strip  $\frac{1}{2} < \sigma_0 \leq 1$ . Then we have*

$$\#\{\chi \mid \text{conductor}(\chi) \text{ a prime} \leq Q \text{ and } L_{f,\chi}(s_0) \overline{L_{g,\chi}(s_0)} \neq 0\} \gg \frac{Q^2}{(\log Q)^4}$$

as  $Q \rightarrow \infty$ . The implied constant depends on  $f, g$  and  $s_0$ .

This theorem should be compared to some non-vanishing results in the theory of automorphic forms. To explain the connection, let  $F$  be a number field,  $S$  be a finite set of places of  $F$ , and  $\pi$  be a unitary cuspidal automorphic representation of  $\operatorname{GL}(n)$  over  $F$ . Let  $s_0 = \sigma_0 + it_0$  be a fixed point in the complex plane. Then Rohrlich [R] proved that for  $n = 1$  and  $2$  there are infinitely many primitive ray class characters  $\chi$  of  $F$  such that  $\chi$  is unramified at the places in  $S$  and  $L(\pi \otimes \chi, s_0) \neq 0$ . For  $n \geq 3$ , Barthel and Ramakrishnan [BR] proved that the same result remains true as long as  $\pi$  is tempered (i.e. satisfies the Ramanujan conjecture) and  $\sigma_0 > 1 - \frac{2}{n+1}$  (see also [LRS] for a related result). For automorphic representations of  $\operatorname{GL}(4)$  over  $\mathbb{Q}$  (the case that is related to this paper) the result of Barthel and Ramakrishnan states that for  $\sigma_0 > \frac{3}{5}$  there are infinitely many primitive Dirichlet characters that  $L(\pi \otimes \chi, s_0) \neq 0$ . Note that our non-vanishing result (Theorem 1.2) surpasses the bound  $\frac{3}{5}$ . This is due to the fact that we are dealing with the product of two

twisted  $\mathrm{GL}(2)$   $L$ -functions  $(L_{f,\chi}(s)\overline{L_{g,\chi}(s)})$  and thus the Gauss sums associated to the functional equations of these two  $L$ -functions cancel each other (see Lemma 2.2). Therefore the contributions from the sums corresponding to  $1 - s_0$  in Lemma 2.2 can be dealt with in ways similar to the sums corresponding to  $s_0$ . This enables us to prove a non-vanishing result in the half plane  $\sigma_0 > \frac{1}{2}$ . In fact a similar result should be true on the line  $\sigma_0 = \frac{1}{2}$ , however establishing such a result needs a more elaborate treatment of the error terms in Proposition 2.5.

In the next two sections we prove the above theorems.

## 2. THEOREM 1.1

Let  $k \geq l$  and  $s_0 = \sigma_0 + it_0$ . We set

$$P_\chi(s_0) = L_{f,\chi}(s_0)\overline{L_{g,\chi}(s_0)}.$$

We first derive an asymptotic formula for  $\sum_{\chi} P_\chi(s_0)$  as  $\chi$  varies over the primitive characters mod  $q$ . Here we do not assume that  $q$  is a prime. Let

$$Z_{s_0}(x) = \frac{1}{2\pi i} \int_{(1)}^{\infty} L_{\infty,k}(s+s_0)L_{\infty,l}(s+\bar{s}_0)x^{-s} \frac{ds}{s}.$$

Writing the integral representations of the  $\Gamma$  functions in the expression for  $Z_{s_0}(x)$  and interchanging the order of integration we arrive at

$$Z_{s_0}(x) = (2\pi)^{-2\sigma_0} \int_0^{\infty} t_1^{\frac{k-1}{2}+s_0-1} e^{-t_1} \left( \int_{\frac{4\pi^2 x}{t_1}}^{\infty} t_2^{\frac{l-1}{2}+\bar{s}_0-1} e^{-t_2} dt_2 \right) dt_1.$$

Note that this representation for  $Z_{s_0}(x)$  shows that  $|Z_{s_0}(x)| \leq Z_{\sigma_0}(x)$ , moreover by integration by parts we can find an expression for  $Z_{s_0}(x)$  in terms of  $K$ -Bessel functions. Applying the standard bounds for  $K$ -Bessel functions yields

$$(2) \quad |Z_{s_0}(x)| \ll \begin{cases} 1 & x \leq 1 \\ x^{\frac{k}{4}+\frac{l}{4}+\sigma_0-\frac{5}{4}} e^{-4\pi\sqrt{x}} & x > 1 \end{cases}$$

(see [A], Lemmas 6.2 and 6.3 for details).

We next represent  $P_\chi(s_0)$  as a sum of two rapidly convergent series.

**Lemma 2.1.** *Let  $f \in S_k(\Gamma_0(M), \psi)$  and  $g \in S_l(\Gamma_0(N), \eta)$  be new forms. Suppose that  $\chi$  is a primitive Dirichlet character modulo  $q$  with  $(q, MN) = 1$ . Then*

$$L_{\infty,k}(s_0)L_{\infty,l}(\bar{s}_0)P_\chi(s_0) = S_{f,g}(s_0) + \epsilon_{f,\chi}\epsilon_{\bar{g},\bar{\chi}}(q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \hat{S}_{f,g}(1-s_0)$$

where

$$S_{f,g}(s_0) = \sum_{m,n \geq 1} \frac{a_f(m)\bar{a}_g(n)}{(mn)^{\sigma_0}} \left(\frac{n}{m}\right)^{it_0} Z_{s_0}\left(\frac{mn}{q^2\sqrt{MN}}\right)\chi(m)\bar{\chi}(n),$$

and

$$\hat{S}_{f,g}(1-s_0) = \sum_{m,n \geq 1} \frac{\bar{a}_f(m)a_g(n)}{(mn)^{1-\sigma_0}} \left(\frac{m}{n}\right)^{it_0} Z_{1-s_0}\left(\frac{mn}{q^2\sqrt{MN}}\right)\bar{\chi}(m)\chi(n).$$

*Proof.* We have

$$S_{f,g}(s_0) = \frac{1}{2\pi i} \int_{(1)} L_{\infty,k}(s+s_0)L_{f,\chi}(s+s_0)L_{\infty,l}(s+\bar{s}_0)L_{\bar{g},\bar{\chi}}(s+\bar{s}_0)(q^2\sqrt{MN})^s \frac{ds}{s}.$$

Moving the line of integration to the left of zero and calculating the residue at  $s = 0$ , and an application of (1) result in

$$\begin{aligned} S_{f,g}(s_0) &= L_{\infty,k}(s_0)L_{\infty,l}(\bar{s}_0)P_{\chi}(s_0) + \epsilon_{f,\chi}\epsilon_{\bar{g},\bar{\chi}}(q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \\ &\times \frac{1}{2\pi i} \int_{(-1)} L_{\infty,k}(1-s-s_0)L_{\bar{f},\bar{\chi}}(1-s-s_0)L_{\infty,l}(1-s-\bar{s}_0)L_{g,\chi}(1-s-\bar{s}_0)(q^2\sqrt{MN})^{-s} \frac{ds}{s}. \end{aligned}$$

Now changing  $s$  to  $-s$  yields the result.  $\square$

From now on for simplicity we let  $L_{\infty}(s_0) = L_{\infty,k}(s_0)L_{\infty,l}(\bar{s}_0)$ . Next we average  $P_{\chi}(s_0)$  over all primitive Dirichlet characters modulo  $q$ . We have

**Lemma 2.2.** *Let  $q \not\equiv 2 \pmod{4}$  and  $(q, MN) = 1$ . Then*

$$L_{\infty}(s_0) \sum_{\chi(\bmod q)}^* P_{\chi}(s_0) = \sum_{d|q} \mu\left(\frac{q}{d}\right)\phi(d) \left( S_{f,g}^d(s_0) + \epsilon_f\epsilon_{\bar{g}}\psi\bar{\eta}(q)(q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \hat{S}_{f,g}^d(1-s_0) \right)$$

where  $\mu$  is the Möbius function,  $\phi$  is the Euler function,

$$S_{f,g}^d(s_0) = \sum_{\substack{m,n, (mn,q)=1 \\ m \equiv n \pmod{d}}} \frac{a_f(m)\bar{a}_g(n)}{(mn)^{\sigma_0}} \left(\frac{n}{m}\right)^{it_0} Z_{s_0}\left(\frac{mn}{q^2\sqrt{MN}}\right),$$

and

$$\hat{S}_{f,g}^d(1-s_0) = \sum_{\substack{m,n, (mn,q)=1 \\ Nm \equiv Mn \pmod{d}}} \frac{\bar{a}_f(m)a_g(n)}{(mn)^{1-\sigma_0}} \left(\frac{m}{n}\right)^{it_0} Z_{1-s_0}\left(\frac{mn}{q^2\sqrt{MN}}\right).$$

*Proof.* From Lemma 2.1 we have

$$L_{\infty}(s_0) \sum_{\chi(\bmod q)}^* P_{\chi}(s_0) = \sum_{\chi(\bmod q)}^* \left( S_{f,g}(s_0) + \epsilon_f\epsilon_{\bar{g}}\psi\bar{\eta}(q)\chi(M)\bar{\chi}(N)(q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \hat{S}_{f,g}(1-s_0) \right).$$

Note that  $\epsilon_{f,\chi}\epsilon_{\bar{g},\bar{\chi}} = \epsilon_f\epsilon_{\bar{g}}\psi\bar{\eta}(q)\chi(M)\bar{\chi}(N)$ . To simplify the above expression, we

need to evaluate  $\sum_{\chi(\bmod q)}^* \chi(m)\bar{\chi}(n)$  and  $\sum_{\chi(\bmod q)}^* \chi(Mn)\bar{\chi}(Nm)$ . Let

$$h_{m,n}(q) = \sum_{\chi(\bmod q)}^* \chi(m)\bar{\chi}(n).$$

We have

$$\sum_{d|q} h_{m,n}(d) = \sum_{\chi(\bmod q)} \chi(m)\bar{\chi}(n) = \begin{cases} \phi(q) & \text{if } m \equiv n \pmod{q} \\ 0 & \text{otherwise.} \end{cases}$$

Now applying the Möbius inversion formula ([M], Section 1.1) on the above identity yields

$$\sum_{\chi(\bmod q)}^* \chi(m)\bar{\chi}(n) = h_{m,n}(q) = \sum_{d|(q,m-n)} \mu\left(\frac{q}{d}\right)\phi(d).$$

Applying this and a similar identity for  $\sum_{\chi(\bmod q)}^* \chi(Mn)\bar{\chi}(Nm)$  in the expression for

$L_\infty(s_0) \sum_{\chi(\bmod q)}^* P_\chi(s_0)$  at the beginning of the proof imply the result.  $\square$

Next we find an asymptotic for the terms in  $S_{f,g}^d(s_0)$  corresponding to  $m = n$ . To explain our result we need to introduce a notation. We know that for any prime  $p$ ,  $a_f(p) = \alpha_{f,1}(p) + \alpha_{f,2}(p)$  and  $a_g(p) = \alpha_{g,1}(p) + \alpha_{g,2}(p)$ , where  $\alpha_{f,1}(p)\alpha_{f,2}(p) = \psi(p)$  and  $\alpha_{g,1}(p)\alpha_{g,2}(p) = \eta(p)$ . Let

$$R_q(s) = \prod_{p|q} \left(1 - \frac{\psi\bar{\eta}(p)}{p^{2s}}\right)^{-1} \prod_{i=1}^2 \prod_{j=1}^2 \left(1 - \frac{\alpha_{f,i}(p)\bar{\alpha}_{g,j}(p)}{p^s}\right).$$

**Lemma 2.3.** *Let  $f, g$  and  $s_0$  be as Theorem 1.1. Then*

$$\sum_{n, (n,q)=1} \frac{a_f(n)\bar{a}_g(n)}{n^{2\sigma_0}} Z_{s_0}\left(\frac{n^2}{q^2\sqrt{MN}}\right) \sim L_\infty(s_0)L(f \otimes g, 2\sigma_0)R_q(2\sigma_0)$$

as  $q \rightarrow \infty$ .

*Proof.* From the definition of  $Z_{s_0}(x)$ , we have

$$\begin{aligned} & \sum_{n, (n,q)=1} \frac{a_f(n)\bar{a}_g(n)}{n^{2\sigma_0}} Z_{s_0}\left(\frac{n^2}{q^2\sqrt{MN}}\right) \\ &= \frac{1}{2\pi i} \int_{(1)} L_{\infty,k}(s+s_0)L_{\infty,l}(s+\bar{s}_0)L(f \otimes g, 2s+2\sigma_0)R_q(2s+2\sigma_0)(q^2\sqrt{MN})^s \frac{ds}{s}. \end{aligned}$$

Moving the line of integration to the left of zero implies the result.  $\square$

The next lemma gives an estimation for the off-diagonal terms in  $S_{f,g}^d(s_0)$ .

**Lemma 2.4.** *Let  $\epsilon > 0$  be arbitrary, then*

$$\sum_{d|q} \mu\left(\frac{q}{d}\right)\phi(d) \sum_{\substack{m,n, (mn,q)=1 \\ m \equiv n \pmod{d}, m \neq n}} \frac{a_f(m)\bar{a}_g(n)}{(mn)^{\sigma_0}} \left(\frac{n}{m}\right)^{it_0} Z_{s_0}\left(\frac{mn}{q^2\sqrt{MN}}\right) = O(q^{2-2\sigma_0+\epsilon}).$$

*The implied constant depends on  $f, g, s_0$  and  $\epsilon$ .*

*Proof.* We closely follow Section 3.2 of [S]. First of all we recall Rankin-Shiu's estimate for the sum of Fourier coefficients of modular forms. Let  $d \neq 1$  and  $(n, d) = 1$ , then for a newform  $g$  we have

$$\sum_{\substack{n \leq x \\ n \equiv m \pmod{d}}} |a_g(n)| \ll \frac{x}{\phi(d)} (\log x)^{-\epsilon_1}$$

as  $x \rightarrow \infty$ , where  $x > d^\alpha$  for  $1 < \alpha < 2$ . Here,  $0 < \epsilon_1 \leq \delta \simeq 0.06\dots$  is arbitrary and the bound is uniform in  $m, d$  and  $\alpha$  (see [S], page 5 for details). We use this together with Rankin's estimate [RA1]

$$\sum_{m \leq x} |a_f(m)| \ll x(\log x)^{-\delta}$$

to bound the inner sum in the statement of the lemma. Let  $1 < \alpha < \frac{10}{9}$  be a fixed number. We only need to find estimates for the following ranges of  $m$  and  $n$ .

- (i)  $n > d^\alpha$ .
- (ii)  $d^{\frac{4}{5}\alpha} \leq n < m \leq d^\alpha$ .
- (iii)  $n < d^{\frac{4}{5}\alpha}$  and  $d \leq m \leq d^\alpha$ .

Now we estimate the inner sum in the statement of the lemma in each case.

(i) We assume  $n > d^\alpha$ . By employing Rankin-Shiu's and Rankin's estimates, bounds for  $Z_{\sigma_0}(x)$  and partial summation we have

$$\sum_{\substack{m \geq \frac{q^2\sqrt{MN}}{d^\alpha} \\ (m,q)=1}} \frac{|a_f(m)|}{m^{\sigma_0}} \sum_{\substack{n > d^\alpha, (n,q)=1 \\ n \equiv m \pmod{d}}} \frac{|a_g(n)|}{n^{\sigma_0}} Z_{\sigma_0}\left(\frac{mn}{q^2\sqrt{MN}}\right) \ll \frac{1}{\phi(d)} (q^2\sqrt{MN})^{1-\sigma_0},$$

and

$$\sum_{\substack{m < \frac{q^2\sqrt{MN}}{d^\alpha} \\ (m,q)=1}} \frac{|a_f(m)|}{m^{\sigma_0}} \sum_{\substack{n > d^\alpha, (n,q)=1 \\ n \equiv m \pmod{d}}} \frac{|a_g(n)|}{n^{\sigma_0}} Z_{\sigma_0}\left(\frac{mn}{q^2\sqrt{MN}}\right) \ll \frac{1}{\phi(d)} (q^2\sqrt{MN})^{1-\sigma_0} (\log(d^{-\alpha} q^2\sqrt{MN}))^{1-\delta}.$$

(ii) Next we consider the range  $d^{\frac{4}{5}\alpha} \leq n < m \leq d^\alpha$ . We recall from the Rankin-Selberg theory [RA2] the asymptotic formula

$$\sum_{n \leq x} |a_g(n)|^2 = c_g x + O(x^{\frac{3}{5}})$$

where  $c_g$  is a constant depending only on  $g$ . By employing Cauchy-Schwarz inequality and above asymptotic we have

$$\sum_{n \leq x} |a_f(n+dt)a_g(n)| \ll x$$

uniformly for  $x \gg (dt)^{\frac{3}{5}}$ . Now by writing  $m = n + dt$  for  $1 \leq t \leq d^{\alpha-1}$ , and applying partial summation, we get the following estimation of the inner sum.

$$\begin{aligned} & \sum_{1 \leq t \leq d^{\alpha-1}} \sum_{d^{\frac{4}{5}\alpha} < n \leq \frac{q^2\sqrt{MN}}{dt}} \frac{|a_f(n+dt)a_g(n)|}{(n+dt)^{\sigma_0} n^{\sigma_0}} Z_{\sigma_0}\left(\frac{(n+dt)n}{q^2\sqrt{MN}}\right) \\ & \ll \sum_{1 \leq t \leq d^{\alpha-1}} \frac{(q^2\sqrt{MN})^{1-\sigma_0}}{dt} \ll \frac{1}{d} (q^2\sqrt{MN})^{1-\sigma_0} \log d. \end{aligned}$$

A similar estimation is true for the range  $\frac{q^2\sqrt{MN}}{dt} < n \leq d^\alpha$ .

(iii) Finally we consider  $m$  and  $n$ 's in the range  $n < d^{\frac{4}{5}\alpha}$  and  $d \leq m \leq d^\alpha$ . Note that since  $1 < \alpha < \frac{10}{9}$  and  $\sigma_0 > \frac{1}{2}$ , we have  $\frac{1}{2} - \frac{9}{10}\alpha(1-\sigma_0) > 0$ . We choose an  $\epsilon$  such that  $0 < \epsilon < \frac{1}{2} - \frac{9}{10}\alpha(1-\sigma_0)$ . By Deligne's bound for the Fourier coefficients of newforms we have

$$|a_f(m)a_g(n)| \ll_\epsilon d^\epsilon.$$

By applying this bound we have

$$\sum_{\substack{n < d^{\frac{4}{5}\alpha}, d \leq m \leq d^\alpha \\ m \equiv n \pmod{d}}} \frac{|a_f(m)a_g(n)|}{(mn)^{\sigma_0}} Z_{\sigma_0}\left(\frac{mn}{q^2\sqrt{MN}}\right) \ll d^{\frac{9}{5}\alpha(1-\sigma_0)-1+\epsilon} \ll d^{-\epsilon}.$$

Applying estimates of (i), (ii) and (iii) to the inner sum in the statement of the lemma implies the result.  $\square$

Combining the results of Lemma 2.2, Lemma 2.3 and Lemma 2.4 we have the following.

**Proposition 2.5.** *Let  $f, g$  and  $s_0$  be as Theorem 1.1 and let  $\epsilon > 0$  be arbitrary. Let  $q \not\equiv 2 \pmod{4}$  and  $(q, MN) = 1$ . We have*

$$\sum_{\chi \pmod{q}}^* P_\chi(s_0) = L(f \otimes g, 2\sigma_0) R_q(2\sigma_0) \sum_{d|q} \mu\left(\frac{q}{d}\right) \phi(d) + O(q^{2-2\sigma_0+\epsilon}).$$

The implied constant depends on  $f, g, s_0$  and  $\epsilon$ .

*Proof.* First of all note that a result similar to Lemma 2.4 is also true for the off-diagonal terms in  $\hat{S}_{f,g}^d(1-s_0)$ , and in this case the corresponding sum is bounded by  $q^{2\sigma_0+\epsilon}$ . For the diagonal terms in  $\hat{S}_{f,g}^d(1-s_0)$  (ones corresponding to  $Nm = Mn$ ), by applying Deligne's bound for Fourier coefficients of new forms we have

$$\begin{aligned} \sum_{\substack{m,n,(mn,q)=1 \\ Nm=Mn}} \frac{|a_f(m)a_g(n)|}{(mn)^{1-\sigma_0}} Z_{1-\sigma_0}\left(\frac{mn}{q^2\sqrt{MN}}\right) &\ll_{M,N} \sum_{n,(n,q)=1} \frac{n^\epsilon}{n^{2(1-\sigma_0)}} Z_{1-\sigma_0}\left(\frac{Mn^2}{q^2N\sqrt{MN}}\right) \\ &\ll q^{2\sigma_0-1+\epsilon}. \end{aligned}$$

Applying these estimates together with Lemmas 2.3 and 2.4 in Lemma 2.2 will imply the desired result.  $\square$

**Proof of Theorem 1.1.** We sum the asymptotic formula given in Lemma 2.5 over primes  $q \leq Q$  where  $(q, MN) = 1$ . Note that for  $q$  prime,  $R_q(2\sigma_0) = 1 + O(q^{-2\sigma_0})$  and  $\sum_{d|q} \mu\left(\frac{q}{d}\right) \phi(d) = q - 2$ . Now the result follows from the prime number theorem.  $\square$

### 3. THEOREM 1.2

First of all in Lemma 2.1 let

$$b_u = \sum_{u=mn} a_f(m) \bar{a}_g(n) \chi(m) \bar{\chi}(n) \left(\frac{n}{m}\right)^{it_0}.$$

Note that by Deligne's bound for Fourier coefficients of newforms we have

$$|b_u| \ll_\epsilon u^\epsilon.$$

So by Lemma 2.1 for given  $X$  we have

$$\begin{aligned} L_\infty(s_0) P_\chi(s_0) &= \sum_{u \leq X} \frac{b_u}{u^{\sigma_0}} Z_{s_0}\left(\frac{u}{q^2\sqrt{MN}}\right) + \sum_{u \geq X} \frac{b_u}{u^{\sigma_0}} Z_{s_0}\left(\frac{u}{q^2\sqrt{MN}}\right) \\ &+ \epsilon_{f,\chi} \epsilon_{\bar{g},\bar{\chi}} (q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \sum_{u \leq X} \frac{\bar{b}_u}{u^{1-\sigma_0}} Z_{1-s_0}\left(\frac{u}{q^2\sqrt{MN}}\right) \\ &+ \epsilon_{f,\chi} \epsilon_{\bar{g},\bar{\chi}} (q^2\sqrt{MN})^{1-2\sigma_0} \left(\frac{N}{M}\right)^{it_0} \sum_{u \geq X} \frac{\bar{b}_u}{u^{1-\sigma_0}} Z_{1-s_0}\left(\frac{u}{q^2\sqrt{MN}}\right) \\ &= L_1(s_0) + L_2(s_0) + L_3(s_0) + L_4(s_0). \end{aligned}$$

From here we have

$$(3) \quad |L_\infty(s_0)|^2 \sum_{\substack{q \leq Q \\ (q, MN)=1}} \frac{q}{\phi(q)} \sum_{\chi \pmod{q}}^* |P_\chi(s_0)|^2 \ll |L_\infty(s_0)|^2 \sum_{i=1}^4 \sum_{q \leq Q} \frac{q}{\phi(q)} \sum_{\chi \pmod{q}}^* |L_i(s_0)|^2.$$

Now let  $X = \sqrt{MN}Q^2(\log Q)^2$ . Then by employing bound (2) for  $Z_{s_0}(x)$ , we have

$$(4) \quad \sum_{q \leq Q} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* |L_i(s_0)|^2 \ll Q^{-19}$$

for  $i = 2, 4$ . We know that by the large sieve inequality for characters we have

$$\sum_{q \leq Q} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* \left| \sum_{u=1}^X a_u \chi(u) \right|^2 \leq (X + 3Q^2) \sum_{u=1}^X |a_u|^2$$

(see [D], page 160, Theorem 4). Let

$$a_u = u^{-\sigma_0} Z_{s_0} \left( \frac{u}{q^2 \sqrt{MN}} \right) \sum_{\substack{u=mn \\ (n,q)=1}} a_f(m) \bar{a}_g(n) \bar{\chi}(n^2) \left( \frac{n}{m} \right)^{it_0}.$$

By Deligne's bound and (2) we have

$$|a_u| \ll_{\epsilon} u^{-\sigma_0 + \epsilon}.$$

This bound together with the large sieve inequality imply that for  $i = 1$

$$(5) \quad \sum_{q \leq Q} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* |L_i(s_0)|^2 = \sum_{q \leq Q} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* \left| \sum_{u=1}^X a_u \chi(u) \right|^2 \ll Q^2 (\log Q)^2.$$

The same bound is valid for  $i = 3$ . Now applying (4) and (5) in (3) implies the following.

**Proposition 3.1.** *Let  $f, g$  and  $s_0$  be as Theorem 1.1. We have*

$$\sum_{\substack{q \leq Q \\ (q, MN)=1}} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* |P_{\chi}(s_0)|^2 \ll Q^2 (\log Q)^2.$$

The implied constant depends on  $f, g$  and  $s_0$ .

**Proof of Theorem 1.2.** By Cauchy-Schwarz inequality we have

$$\#\{\chi \mid \text{conductor}(\chi) \text{ a prime} \leq Q \text{ and } P_{\chi}(s_0) \neq 0\} \geq \frac{\left| \sum_{\substack{q \leq Q, q \text{ prime} \\ (q, MN)=1}} \sum_{\chi(\bmod q)}^* P_{\chi}(s_0) \right|^2}{\sum_{\substack{q \leq Q \\ (q, MN)=1}} \frac{q}{\phi(q)} \sum_{\chi(\bmod q)}^* |P_{\chi}(s_0)|^2}.$$

The result follows from this inequality, Theorem 1.1 and Proposition 3.1.  $\square$

#### ACKNOWLEDGMENT

The author would like to thank the referee for helpful comments and suggestions. I also would like to thank Hershy Kisilevsky for several helpful discussions related to this work.

## REFERENCES

- [A] A. Akbary, Non-vanishing of weight  $k$  modular  $L$ -functions with large level, *J. Ramanujan Math. Soc.* **14** (1999), 37–54.
- [BR] L. Barthel and D. Ramakrishnan, A nonvanishing result for twists of  $L$ -functions of  $GL(n)$ , *Duke Math. J.* **74** (1994), 681–700.
- [D] H. Davenport, *Multiplicative Number Theory*, third edition, Springer, 2000.
- [LRS] W. Luo, Z. Rudnick, and P. Sarnak, On the generalized Ramanujan conjecture for  $GL(n)$ , In Automorphic forms, automorphic representations, and arithmetic (Fort Worth, TX, 1996), 301–310, *Proc. Sympos. Pure Math.*, **66**, Part 2, AMS, 1999.
- [E] P. D. T. A. Elliott, On the distribution of the values of Dirichlet  $L$ -series in the half plane  $\sigma > \frac{1}{2}$ , *Indag. Math.* **33** (1971), 222–234.
- [M] M. R. Murty, *Problems in Analytic Number Theory*, Springer, 2001.
- [R] D. Rohrlich, Nonvanishing of  $L$ -functions for  $GL(2)$ , *Invent. Math.* **97** (1989), 381–403.
- [RA1] R. A. Rankin, Sums of powers of cusp form coefficients, II, *Math. Ann.* **272** (1985), 593–600.
- [RA2] R. A. Rankin, Contributions to the theory of Ramanujan’s function  $\tau(n)$  and similar arithmetical functions. II. The order of the Fourier coefficients of integral modular forms, *Proc. Cambridge Phil. Soc.* **35** (1939), 357–372.
- [S] T. Stefanicki, Non-vanishing of  $L$ -functions attached to automorphic representations of  $GL(2)$  over  $Q$ , *J. reine angew. Math.* **474** (1996), 1–24.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF LETHBRIDGE, 4401  
UNIVERSITY DRIVE WEST, LETHBRIDGE, ALBERTA, T1K 3M4, CANADA  
*E-mail address:* akbary@cs.uleth.ca